

Product datasheet

Anti-SLC4A4 antibody ab56512

画像数 1

製品の概要

製品名	Anti-SLC4A4 antibody
製品の詳細	Mouse monoclonal to SLC4A4
由来種	Mouse
アプリケーション	適用あり: WB
種交差性	交差種: Human
免疫原	Recombinant full length protein, corresponding to amino acids 1-647 of Human SLC4A4
特記事項	Abcam is committed to meeting high standards of ethical manufacturing and as such, we will be discontinuing this product, which has been generated by the ascites method, within the next year. We are sorry for any inconvenience this may cause. If you would like help finding an alternative product, please do not hesitate to contact our scientific support team.

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
バッファー	Preservative: None PBS, pH 7.2
精製度	Protein G purified
ポリ/モノ	モノクローナル
アイソタイプ	IgG2a
軽鎖の種類	kappa

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab56512** in the following tested applications.


The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	Abreviews	特記事項
WB		<p>Use a concentration of 1 - 5 µg/ml.</p> <p>This antibody has only been tested in WB against the recombinant fragment used as immunogen. We have no data on the detection of endogenous protein.</p>

ターゲット情報

機能	Electrogenic sodium/bicarbonate cotransporter with a Na(+):HCO ₃ (-) stoichiometry varying from 1:2 to 1:3. May regulate bicarbonate influx/efflux at the basolateral membrane of cells and regulate intracellular pH.
組織特異性	Isoform 1 is expressed in pancreas and to a lower extent in heart, skeletal muscle, liver, parotid salivary glands, prostate, colon, stomach, thyroid, brain and spinal chord. Corneal endothelium cells express only isoform 1 (at protein level). Isoform 2 is specifically expressed in kidney at the level of proximal tubules.
関連疾患	<p>Defects in SLC4A4 are the cause of proximal renal tubular acidosis with ocular abnormalities (pRTA-OA) [MIM:604278]; also known as renal tubular acidosis II. Caused by an impairment of bicarbonate absorption in the proximal tubule, proximal renal tubular acidosis (pRTA) is characterized by a decreased renal HCO₃(-) threshold. pRTA-OA is an extremely rare autosomal recessive syndrome characterized by short stature, profound pRTA, mental retardation, bilateral glaucoma, cataracts and bandkeratopathy.</p> <p>Note=Loss of interaction with and stimulation by CA4 is the cause of retinitis pigmentosa type 17 (RP17).</p>
配列類似性	Belongs to the anion exchanger (TC 2.A.31) family.
翻訳後修飾	<p>Phosphorylation of Ser-1026 by PKA increases the binding of CA2 and changes the Na(+):HCO₃(-) stoichiometry of the transporter from 3:1 to 2:1. Phosphorylation of Thr-49 regulates isoform 1 conductance.</p> <p>N-glycosylation is not necessary for the transporter basic functions.</p>
細胞内局在	Basolateral cell membrane.

画像



Western blot against tagged recombinant protein immunogen using ab56512 SLC4A4 antibody at 1µg/ml. Predicted band size of immunogen is 97 kDa

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors