

Product datasheet

Anti-UCP1 antibody ab57687

画像数 2

製品の概要

製品名	Anti-UCP1 antibody
製品の詳細	Mouse monoclonal to UCP1
アプリケーション	適用あり: WB, Flow Cyt
種交差性	交差種: Human
免疫原	Recombinant fragment: PVDVVKTRFI NSPPGQYKSV PNCAMKVFTN EGPTAF, corresponding to amino acids 232-268 of Human UCP1 Run BLAST with ExPASy Run BLAST with NCBI

特記事項

Abcam is committed to meeting high standards of ethical manufacturing and has decided to discontinue this product by June 2019 as it has been generated by the ascites method. We are sorry for any inconvenience this may cause.

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
バッファー	Preservative: None PBS, pH 7.2
精製度	Protein G purified
ポリモノ	モノクローナル
アイソタイプ	IgG2a
軽鎖の種類	kappa

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab57687** in the following tested applications.


The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	Abreviews	特記事項
WB		<p>Use a concentration of 1 - 5 µg/ml.</p> <p>This antibody has only been tested in WB against the recombinant fragment used as immunogen. We have no data on the detection of endogenous protein.</p>
Flow Cyt		<p>Use 0.1 µg for 10⁶ cells. ab170191-Mouse monoclonal IgG2a, is suitable for use as an isotype control with this antibody.</p>

ターゲット情報


機能	UCP are mitochondrial transporter proteins that create proton leaks across the inner mitochondrial membrane, thus uncoupling oxidative phosphorylation from ATP synthesis. As a result, energy is dissipated in the form of heat.
組織特異性	Brown adipose tissue.
配列類似性	<p>Belongs to the mitochondrial carrier family.</p> <p>Contains 3 Solcar repeats.</p>
細胞内局在	Mitochondrion inner membrane.
製品の状態	UCP1 is preferentially expressed in brown adipose tissue

画像



Flow Cytometry - Anti-UCP1 antibody (ab57687)

Overlay histogram showing HepG2 cells stained with ab57687 (red line). The cells were fixed with 4% paraformaldehyde (10 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab57687, 0.1µg/1x10⁶ cells) for 30 min at 22°C. The secondary antibody used was an anti-mouse Alexa Fluor® 488 (ab150113) at 1/2000 dilution for 30 min at 22°C. Isotype control antibody (black line) was mouse IgG2a [ICIGG2A] (ab91361, 0.1µg/1x10⁶ cells) used under the same conditions. Unlabelled sample (blue line) was also used as a control. Acquisition of >5,000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter. This antibody gave a positive signal in HepG2 cells fixed with 80% methanol (5 min)/permeabilized with 0.1% PBS-Tween for 20 min used under the same conditions.



Western blot - UCP1 antibody (ab57687)

Western blot against tagged recombinant protein immunogen using ab57687 UCP1 antibody at 1µg/ml. Predicted band size of immunogen is 30 kDa

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish

- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors