

Product datasheet

Anti-p73 antibody ab70093

画像数 2

製品の概要

製品名	Anti-p73 antibody
製品の詳細	Rabbit polyclonal to p73
由来種	Rabbit
特異性	ab70093 detects endogenous levels of total p73 protein.
アプリケーション	適用あり: WB, ELISA, IHC-P
種交差性	交差種: Human
免疫原	Synthetic peptide derived from an internal sequence of human p73.
ポジティブ・コントロール	WB: Jurkat cell extract IHC-P: Human brain tissue

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
バッファー	Preservative: 0.02% Sodium Azide Constituents: 50% Glycerol, PBS (without Mg ²⁺ and Ca ²⁺), 150mM Sodium chloride, pH 7.4
精製度	Immunogen affinity purified
特記事項 (精製)	ab70093 was affinity purified from rabbit antiserum by affinity chromatography using epitope specific immunogen.
ポリモノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab70093** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


アプリケーション	Abreviews	特記事項
WB		1/500 - 1/1000. Detects a band of approximately 70 kDa (predicted molecular weight: 70 kDa).

アプリケーション	Abreviews	特記事項
ELISA		1/40000.
IHC-P		Use at an assay dependent concentration.


ターゲット情報

機能	Participates in the apoptotic response to DNA damage. Isoforms containing the transactivation domain are pro-apoptotic, isoforms lacking the domain are anti-apoptotic and block the function of p53 and transactivating p73 isoforms. May be a tumor suppressor protein.
組織特異性	Expressed in striatal neurons of patients with Huntington disease (at protein level). Brain, kidney, placenta, colon, heart, liver, spleen, skeletal muscle, prostate, thymus and pancreas. Highly expressed in fetal tissue.
配列類似性	Belongs to the p53 family. Contains 1 SAM (sterile alpha motif) domain.
ドメイン	Possesses an acidic transactivation domain, a central DNA binding domain and a C-terminal oligomerization domain that binds to the ABL tyrosine kinase SH3 domain. The WW-binding motif mediates interaction with WWOX.
翻訳後修飾	Isoform alpha (but not isoform beta) is sumoylated on Lys-627, which potentiates proteasomal degradation but does not affect transcriptional activity. Higher levels of phosphorylation seen in the brain from patients with Huntington disease. Ubiquitinated; leading to its degradation by the proteasome.
細胞内局在	Nucleus. Accumulates in the nucleus in response to DNA damage.

画像



ab70093, at 1/50 dilution, staining p73 in paraffin embedded human brain tissue by Immunohistochemistry in the absence or presence of the immunising peptide.



All lanes : Anti-p73 antibody (ab70093) at 1/500 dilution

Lane 1 : Jurkat cell extract

Lane 2 : Jurkat cell extract with immunising peptide at 5 µg

Lysates/proteins at 5 µg per lane.

Predicted band size: 70 kDa

Observed band size: 70 kDa

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors