

Product datasheet

Anti-CD79a antibody [HM47/A9] (Biotin) ab79437

1 References

製品の概要

製品名	Anti-CD79a antibody [HM47/A9] (Biotin)
製品の詳細	Mouse monoclonal [HM47/A9] to CD79a (Biotin)
由来種	Mouse
標識	Biotin
特異性	We have data to indicate that this antibody may not cross react with Opossum. However, this has not been conclusively tested and expression levels may vary in certain cell lines/tissues.
アプリケーション	適用あり: WB
種交差性	交差種: Mouse, Rat, Rabbit, Horse, Chicken, Guinea pig, Cow, Human, Pig, Monkey
免疫原	Synthetic peptide: GTYQDVGSLNIADVQ , corresponding to amino acids 208-222 of human CD79a <div style="text-align: right;"> Run BLAST with Run BLAST with </div>
エピトープ	GTYQDVGSLNIADVQ
ポジティブ・コントロール	Raji cells. Lymph node or tonsil.

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	Preservative: 0.09% Sodium Azide Constituents: 0.2% BSA, 10mM PBS, pH 7.4
精製度	Protein G purified
特記事項(精製)	ab79437 was purified from ascites fluid by Protein G chromatography.
ポリ/モノ	モノクローナル
クローン名	HM47/A9
アイソタイプ	IgG1
軽鎖の種類	kappa

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab79437** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	Abreviews	特記事項
WB		Use a concentration of 0.5 - 1 µg/ml. Predicted molecular weight: 25 kDa.

ターゲット情報

機能	Required in cooperation with CD79B for initiation of the signal transduction cascade activated by binding of antigen to the B-cell antigen receptor complex (BCR) which leads to internalization of the complex, trafficking to late endosomes and antigen presentation. Also required for BCR surface expression and for efficient differentiation of pro- and pre-B-cells. Stimulates SYK autophosphorylation and activation. Binds to BLNK, bringing BLNK into proximity with SYK and allowing SYK to phosphorylate BLNK. Also interacts with and increases activity of some Src-family tyrosine kinases. Represses BCR signaling during development of immature B cells.
組織特異性	B-cells.
関連疾患	Defects in CD79A are the cause of agammaglobulinemia type 3 (AGM3) [MIM:613501]. It is a primary immunodeficiency characterized by profoundly low or absent serum antibodies and low or absent circulating B cells due to an early block of B-cell development. Affected individuals develop severe infections in the first years of life. Note=Two different mutations, one at the splice donor site of intron 2 and the other at the splice acceptor site for exon 3, have been identified. Both mutations give rise to a truncated protein.
配列類似性	Contains 1 Ig-like C2-type (immunoglobulin-like) domain. Contains 1 ITAM domain.
翻訳後修飾	Phosphorylated on tyrosine, serine and threonine residues upon B-cell activation. Phosphorylation of tyrosine residues by Src-family kinases is an early and essential feature of the BCR signaling cascade. The phosphorylated tyrosines serve as docking sites for SH2-domain containing kinases, leading to their activation which in turn leads to phosphorylation of downstream targets. Phosphorylation of serine and threonine residues may prevent subsequent tyrosine phosphorylation.
細胞内局在	Cell membrane. Following antigen binding, the BCR has been shown to translocate from detergent-soluble regions of the cell membrane to lipid rafts although signal transduction through the complex can also occur outside lipid rafts.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you

- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors