

# Anti-Urokinase antibody [PGM2001] ab8475

★★★★★ 2 Abreviews

## 製品の概要

<b>製品名</b>	Anti-Urokinase antibody [PGM2001]
<b>製品の詳細</b>	Mouse monoclonal [PGM2001] to Urokinase
<b>特異性</b>	This antibody is highly specific for urokinase. No cross reactions were found with other serine proteases (tissue plasminogen activator, plasmin, trypsin, chymotrypsin, elastase, thrombin). This antibody binds to human high molecular weight single-chain urokinase type plasminogen activator (HMW-scuPA, 54 kDa), to high molecular weight two-chain urokinase type plasminogen activator (HMW-tcuPA, 52 kDa) and to low molecular weight single-chain urokinase type plasminogen activator (LMW-scuPA, 33 kDa) with high affinity (Kd= 0.1 nM) It does not inhibit the ability of urokinase to activate plasminogen. Urokinase reaction with synthetic substrate S2444 is not inhibited. Human whole plasma or plasma proteins do not interfere with this antibodies binding to urokinase.
<b>アプリケーション</b>	<b>適用あり:</b> Other, Dot blot, ELISA, IHC-Fr, RIA, WB
<b>種交差性</b>	<b>交差種:</b> Human
<b>免疫原</b>	Urinary Urokinase (Human).
<b>特記事項</b>	Method of purification: PEG6000 precipitation + DEAE cellulose chromatography, gradient elution. The buffer composition before lyophilisation is PBS with manitol. After reconstitution the concentration is 0.2 mg/mL.

## 製品の特性

<b>製品の状態</b>	Lyophilised:Please reconstitute with 1ml sterile distilled water and add preservative if required.
<b>保存方法</b>	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term.
<b>バッファー</b>	Lyophilized purified ascites
<b>精製度</b>	Ascites
<b>特記事項(精製)</b>	PEG 6000 precipitation + DEAE-cellulose chromatography, gradient elution
<b>ポリモノ</b>	モノクローナル
<b>クローン名</b>	PGM2001
<b>アイソタイプ</b>	IgG1

## アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab8475** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリ ケー ション	Abreviews	特記事項
Other		Use at an assay dependent concentration.
Dot blot		Use at an assay dependent concentration.
AP		Use at an assay dependent concentration.
ELISA		1/5000. This antibody can be used for quantitative, specific and sensitive assay for urokinase in human fluids (RIA, ELISA) in combination with the non-inhibitory antibody <a href="#">ab8473</a> . This antibody is very stable and can, after appropriate binding, be used over 20-40 times for immunopurification of urokinase in 1 M guanidine chloride.
IHC- Fr		1/30 - 1/50.
RIA		Use at an assay dependent concentration.
WB	★★★★★	Use a concentration of 1 - 3 µg/ml.

## ターゲット情報

機能	Specifically cleave the zymogen plasminogen to form the active enzyme plasmin.
組織特異性	Expressed in the prostate gland and prostate cancers.
配列類似性	Belongs to the peptidase S1 family. Contains 1 EGF-like domain. Contains 1 kringle domain. Contains 1 peptidase S1 domain.
翻訳後修飾	Phosphorylation of Ser-158 and Ser-323 abolishes proadhesive ability but does not interfere with receptor binding.
細胞内局在	Secreted.

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

## Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors