

Product datasheet

Anti-C4 binding protein antibody ab8788

★★★★★ 2 Abreviews 4 References 画像数 1

製品の概要

<b>製品名</b>	Anti-C4 binding protein antibody
<b>製品の詳細</b>	Sheep polyclonal to C4 binding protein
<b>特異性</b>	This product gives a single arc when tested by IEP against fresh human plasma. Identity has been confirmed by double diffusion (Ouchterlony) vs fresh human plasma and a known anti human C4BP.
<b>アプリケーション</b>	<b>適用あり:</b> WB, Flow Cyt, Double Immunodiffusion, Immunoelectrophoresis, Other
<b>種交差性</b>	<b>交差種:</b> Human
<b>免疫原</b>	Human C4BP purified from plasma and shown to be homogeneous by SDS-PAGE.
<b>特記事項</b>	C4-binding protein (C4BP) is a multimeric protein, synthesised in the liver. The most abundant form consists of seven alpha-chains (70kDa) and one beta-chain (45kDa) all linked by disulphide bonds to form the native protein with a molecular weight of 570kDa. It is a plasma protein with a concentration of about 150mg/L in normal plasma. It is a component of the classical complement pathway. C4BP down regulates complement activity in two ways, it binds to C4b thus inhibiting the formation of C3-convertase (C4bC3a) and it accelerates the decay of existing convertases. C4BP acts as a cofactor in Factor I mediated C4b proteolysis. C4BP may also have a regulatory role in the coagulation system, by binding to Protein S. When bound to C4BP, Protein S is inactive.

製品の特性

<b>製品の状態</b>	Liquid
<b>保存方法</b>	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C or -80°C. Avoid freeze / thaw cycle.
<b>バッファー</b>	Preservative: 0.1% Sodium Azide Constituents: Glycine buffered saline, 0.1% EACA, 0.01% Benzamidine, 1mM EDTA, pH 7.4
<b>精製度</b>	IgG fraction
<b>特記事項 (精製)</b>	Antiserum is prepared by immunisation of sheep with human C4BP and, if necessary, adsorbed to monospecificity by use of solid-phase adsorbents. An immunoglobulin fraction is then produced. The titre is adjusted so that inter-batch variation is within 10%. The product is 0.2µm filtered.
<b>一次抗体 備考</b>	C4-binding protein (C4BP) is a multimeric protein, synthesised in the liver. The most abundant

form consists of seven alpha-chains (70kDa) and one beta-chain (45kDa) all linked by disulphide bonds to form the native protein with a molecular weight of 570kDa. It is a plasma protein with a concentration of about 150mg/L in normal plasma. It is a component of the classical complement pathway. C4BP down regulates complement activity in two ways, it binds to C4b thus inhibiting the formation of C3-convertase (C4bC3a) and it accelerates the decay of existing convertases. C4BP acts as a cofactor in Factor I mediated C4b proteolysis. C4BP may also have a regulatory role in the coagulation system, by binding to Protein S. When bound to C4BP, Protein S is inactive.

ポリ/モノ	ポリクローナル
アイソタイプ	IgG
軽鎖の種類	unknown

## アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab8788** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	Abreviews	特記事項
WB	★★★★☆	Use at an assay dependent concentration.
Flow Cyt	★★★★★	Use at an assay dependent concentration. <a href="#">ab37385</a> -Sheep polyclonal IgG, is suitable for use as an isotype control with this antibody.
Double Immunodiffusion		Use at an assay dependent concentration.
Immunoelectrophoresis		Use at an assay dependent concentration.
Other		Use at an assay dependent concentration.

## ターゲット情報

機能	Controls the classical pathway of complement activation. It binds as a cofactor to C3b/C4b inactivator (C3bINA), which then hydrolyzes the complement fragment C4b. It also accelerates the degradation of the C4bC2a complex (C3 convertase) by dissociating the complement fragment C2a. It also interacts with anticoagulant protein S and with serum amyloid P component. The beta chain binds protein S.
配列類似性	Contains 3 Sushi (CCP/SCR) domains.
細胞内局在	Secreted.

## 画像




Image from Pham CT et al, *J Biol Chem.*  
 2011 Jan 7;286(1):123-30. Epub 2010 Nov 3,  
 Fig 8. DOI 10.1074/jbc.M110.180760

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors