

Product datasheet

Anti-TFIISH antibody ab88595

画像数 3

製品の概要

| | |
|--------------|---|
| 製品名 | Anti-TFIISH antibody |
| 製品の詳細 | Mouse polyclonal to TFIISH |
| 由来種 | Mouse |
| アプリケーション | 適用あり: WB, ICC/IF |
| 種交差性 | 交差種: Human |
| 免疫原 | Full length Human TFIISH (NP_003187.1) |
| ポジティブ・コントロール | Human placenta tissue lysate, TFIISH transfected 293T cell lysate, HepG2 cells. |

製品の特性

| | |
|--------|---|
| 製品の状態 | Liquid |
| 保存方法 | Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles. |
| バッファー | Preservative: None Constituents: 1X PBS, pH 7.2 |
| 精製度 | Protein A purified |
| ポリ/モノ | ポリクローナル |
| アイソタイプ | IgG |

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab88595** in the following tested applications.


The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| アプリケーション | Abreviews | 特記事項 |
|----------|-----------|--|
| WB | | Use a concentration of 1 µg/ml. Detects a band of approximately 39 kDa (predicted molecular weight: 39 kDa). |
| ICC/IF | | Use a concentration of 10 µg/ml. |

ターゲット情報

| | |
|-------|--|
| 機能 | Necessary for efficient RNA polymerase II transcription elongation past template-encoded arresting sites. The arresting sites in DNA have the property of trapping a certain fraction of elongating RNA polymerases that pass through, resulting in locked ternary complexes. Cleavage of the nascent transcript by S-II allows the resumption of elongation from the new 3'-terminus. |
| 配列類似性 | Belongs to the TFS-II family. Contains 1 TFIIIS central domain. Contains 1 TFIIIS N-terminal domain. Contains 1 TFIIIS-type zinc finger. |
| 細胞内局在 | Nucleus. |


画像



Anti-TFIIISH antibody (ab88595) at 1 μ g/ml +
Human placenta tissue lysate at 50 μ g

Predicted band size: 39 kDa

Observed band size: 39 kDa



All lanes : Anti-TFIIISH antibody (ab88595) at
1 μ g/ml


Lane 1 : TFIIISH transfected 293T cell lysate

Lane 2 : Non-transfected lysate

Lysates/proteins at 25 μ g per lane.

Predicted band size: 39 kDa

Observed band size: 39 kDa



ab88595 at 10 μ g/ml staining TFIIISH in HepG2 cells by Immunofluorescence.

Immunocytochemistry/ Immunofluorescence -
TFIIISH antibody (ab88595)

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors