


Product datasheet

Anti-RGS9 antibody ab97702

画像数 1

製品の概要

製品名	Anti-RGS9 antibody
製品の詳細	Rabbit polyclonal to RGS9
由来種	Rabbit
アプリケーション	適用あり: WB
種交差性	交差種: Human 交差が予測される動物種: Rat, Horse, Guinea pig, Cat 
免疫原	Synthetic peptide corresponding to a region within internal sequence amino acids 624-673 of Human RGS9 (NP_003826) Run BLAST with ExPASy Run BLAST with NCBI
ポジティブ・コントロール	Human fetal spleen lysate

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
バッファー	Preservative: None Constituents: 2% Sucrose, PBS
精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab97702** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


アブリケーション	Abreviews	特記事項
----------	-----------	------

WB Use a concentration of 1 µg/ml. Predicted molecular weight: 77 kDa. Optimal concentration of antibody depends the protein level in a given sample. Our recommended concentrations are the starting point for general use. Customers are advised have to do a series of titration tests to determine optimal concentration. Good results were obtained when blocked with 5% non-fat dry milk in 0.05% PBS-T.

ターゲット情報

機能	Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form. Binds to G(t)-alpha. Involved in phototransduction; key element in the recovery phase of visual transduction.
組織特異性	Highly expressed in the caudate and putamen, lower levels found in the hypothalamus and nucleus accumbens and very low levels in cerebellum. Not expressed in globus pallidus or cingulate cortex. Isoform 2 is expressed predominantly in pineal gland and retina. Isoform 3 is expressed in retina (abundant in photoreceptors).
関連疾患	Defects in RGS9 are a cause of prolonged electroretinal response suppression (PERRS) [MIM:608415]; also known as bradyopsia. PERRS is characterized by difficulty adjusting to sudden changes in luminance levels mediated by cones.
配列類似性	Contains 1 DEP domain. Contains 1 G protein gamma domain. Contains 1 RGS domain.
ドメイン	In photoreceptor cells the DEP domain is essential for targeting RGS9 to the outer rod segments.
翻訳後修飾	Retinal isoform 3 is light-dependent phosphorylated at 'Ser-478'. Phosphorylation is decreased by light exposition.
細胞内局在	Membrane. Isoform 3 is targeted to the membrane via its interaction with RGS9BP.

画像



Anti-RGS9 antibody (ab97702) at 1 µg/ml (in 5% skim milk / PBS buffer) + human fetal spleen lysate at 10 µg

Secondary

HRP conjugated anti-Rabbit IgG at 1/50000 dilution

Predicted band size: 77 kDa

12% Gel concentration

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors