

Product datasheet

Anti-alpha+beta SNAP antibody [6B7-3] ab69742

製品の概要

| | |
|--------------|---|
| 製品名 | Anti-alpha+beta SNAP antibody [6B7-3] |
| 製品の詳細 | Mouse monoclonal [6B7-3] to alpha+beta SNAP |
| 由来種 | Mouse |
| 特異性 | This antibody detects a doublet of 35 and 36 kDa proteins, corresponding to the apparent molecular mass of alpha + beta SNAP. |
| アプリケーション | 適用あり: WB, IP |
| 種交差性 | 交差種: Mouse, Rat, Hamster, Cow, Dog, Human |
| 免疫原 | Recombinant full length bovine alpha SNAP protein expressed in E coli |
| ポジティブ・コントロール | Rat brain tissue extract and mouse brain tissue extract. |

製品の特性

| | |
|--------|---|
| 製品の状態 | Liquid |
| 保存方法 | Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles. |
| バッファー | Preservative: None Constituents: 50% Glycerol, PBS, 0.1mM PMSF, pH 7.2 |
| 精製度 | Protein G purified |
| ポリ/モノ | モノクローナル |
| クローン名 | 6B7-3 |
| アイソタイプ | IgG1 |
| 軽鎖の種類 | kappa |

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab69742** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| アプリケーション | Abreviews | 特記事項 |
|----------|-----------|---|
| WB | | Use a concentration of 1.5 µg/ml. Predicted molecular weight: 38 kDa. |

| アプリケーション | Abreviews | 特記事項 |
|----------|-----------|-------------------------------------|
| IP | | Use at an assay dependent dilution. |

ターゲット情報

| | |
|--------------|---|
| 関連性 | Alpha, beta and gamma SNAPs, also known as soluble NSF attachment proteins, have apparent molecular weights on SDS PAGE of 35, 36 and 39 kDa, respectively. SNAP proteins mediate the association of NSF to membranes during vesicle fusion reaction, and play a prominent role in synaptic vesicle fusion. In addition, beta SNAP interacts with synaptotagmin and may be involved in calcium regulated exocytosis. SNAP was identified as being required for vesicle fusion in intra Golgi transport reaction in vitro. Alpha SNAP mediates the association of NSF to membranes during vesicle fusion reaction, and plays a prominent role and in synaptic vesicle fusion. Together with synaptotagmin and NSF, alpha SNAP modulates the interaction of SNAP25, VAMP, and syntaxin in, an ATP dependent manner, to form a complex with a sedimentation coefficient of 20 S. This complex may represent an intermediate involved in synaptic vesicle docking and fusion. |
| 細胞内局在 | Cytoplasmic peripheral membrane protein. |

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors