

## Product datasheet

# Anti-eIF3K antibody ab85968

### 画像数 2

#### 製品の概要

製品名	Anti-eIF3K antibody
製品の詳細	Rabbit polyclonal to eIF3K
由来種	Rabbit
アプリケーション	適用あり: WB, IP
種交差性	交差種: Human 交差が予測される動物種: Mouse, Rat, Rabbit, Horse, Guinea pig, Cow, Cat, Dog, Pig, Chimpanzee, Rhesus monkey, Gorilla
免疫原	Synthetic peptide designed within residues: SQLKVVMSKY GWSADESGQI FICSQEE SIKPKNVE KIDFDSVSSI MASSQ, corresponding to C terminal amino acids 168-218 of Human eIF3K (NP_037366.1) <a href="#">Run BLAST with ExPASy</a> <a href="#">Run BLAST with NCBI</a>
ポジティブ・コントロール	Whole cell lysates from HeLa or 293T cells.

#### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
バッファー	Preservative: 0.09% Sodium Azide Constituents: 0.1% BSA, Tris buffered saline
精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

#### アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab85968** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	Abreviews	特記事項
WB		1/2000 - 1/10000. Predicted molecular weight: 25 kDa.

アプリケーション

Abreviews

特記事項

IP

Use at 2-5 µg/mg of lysate.

ターゲット情報

機能

Component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis. The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNA<sub>i</sub> and eIF-5 to form the 43S preinitiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation.

組織特異性

Ubiquitous, with the highest levels of expression in brain, testis and kidney.


配列類似性

Belongs to the eIF-3 subunit K family.

細胞内局在

Nucleus. Cytoplasm.

画像



**All lanes :** Anti-eIF3K antibody (ab85968) at 0.04 µg/ml

**Lane 1 :** HeLa whole cell lysate at 50 µg


**Lane 2 :** HeLa whole cell lysate at 15 µg

**Lane 3 :** HeLa whole cell lysate at 5 µg

**Lane 4 :** 293T whole cell lysate at 50 µg

**Predicted band size:** 25 kDa

**Exposure time:** 10 seconds



1 mg of HeLa whole cell lysate was

immunoprecipitated using 3µg ab85968.

Subsequently, 20% of IP was loaded per lane and probed with ab85968 at 1µg/ml (lane 1) or with a control IgG (lane 2).

Detection: chemiluminescence with exposure time of 3 seconds.

### **Our Abpromise to you: Quality guaranteed and expert technical support**

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors