


# Betulinic acid, TGR5 agonist ab120654

**1 References** [画像数 2](#)


### 製品の概要

製品名	Betulinic acid, TGR5 agonist
製品の詳細	Anticancer agent. TGR5 agonist.
生理活性の詳細	Triterpenoid with various biological activities. Apoptosis inducer, with anticancer properties. Also shows anti-HIV, antimalarial and anti-inflammatory activity. TGR5 agonist (EC <sub>50</sub> = 1.04 μM).
CAS 番号	472-15-1
構造式	




### 製品の特性

体系名	(3β)-3-Hydroxylup-20(29)-en-28-oic acid
分子量	456.70
分子式	C <sub>30</sub> H <sub>48</sub> O <sub>3</sub>
PubChem 登録番号	64971
保存方法	Store at +4°C. The product can be stored for up to 12 months.
溶解性	Soluble in DMSO to 50 mM (with warming)
使用に関する注意	<p>Wherever possible, you should prepare and use solutions on the same day. However, if you need to make up stock solutions in advance, we recommend that you store the solution as aliquots in tightly sealed vials at -20°C. Generally, these will be useable for up to one month. Before use, and prior to opening the vial we recommend that you allow your product to equilibrate to room temperature for at least 1 hour.</p> <p>Need more advice on solubility, usage and handling? Please visit our <a href="#">frequently asked questions (FAQ) page</a> for more details.</p>
由来	Synthetic



Chemical Structure - Betulinic acid, TGR5 agonist  
(ab120654)

2D chemical structure image of ab120654, Betulinic acid, TGR5 agonist



Functional Studies - Betulinic acid, TGR5 agonist  
(ab120654)

**ab32557** staining p38 (phospho T180 + Y182) in MALME-3M cells treated with betulinic acid (ab120654), by ICC/IF. Increase in expression of p38 (phospho T180 + Y182) correlates with increased concentration of betulinic acid, as described in literature. The cells were incubated at 37°C for 30 minutes in media containing different concentrations of ab120654 (betulinic acid) in DMSO, fixed with 4% formaldehyde for 10 minutes at room temperature and blocked with PBS containing 10% goat serum, 0.3 M glycine, 1% BSA and 0.1% tween for 2h at room temperature. Staining of the treated cells with **ab32557** (1/200 dilution) was performed overnight at 4°C in PBS containing 1% BSA and 0.1% tween. A DyLight 488 goat anti-rabbit polyclonal antibody (**ab96899**) at 1/250 dilution was used as the secondary antibody. Nuclei were counterstained with DAPI and are shown in blue.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES, NOT FOR USE IN HUMANS"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise,

please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

## Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors
- Abcam biochemicals are novel compounds and we have not tested their biological activity in house. Please use the literature to identify how to use these products effectively. If you require further assistance please contact the scientific support team