


CCCP, Mitochondrial oxidative phosphorylation uncoupler ab141229

16 References 画像数 5

医薬用外劇物

製品の概要

製品名	CCCP, Mitochondrial oxidative phosphorylation uncoupler
製品の詳細	Potent mitochondrial oxidative phosphorylation uncoupler
生理活性の詳細	Potent mitochondrial oxidative phosphorylation uncoupler. Renders mitochondrial inner membrane permeable to protons. Induces apoptosis <i>in vitro</i> .
精製度	> 99%
CAS 番号	555-60-2
構造式	

製品の特性

体系名	2-[2-(3-Chlorophenyl)hydrazinylidene]propanedinitrile
分子量	204.62
分子式	C ₉ H ₅ ClN ₄
PubChem 登録番号	2603
保存方法	Store at -20°C. Store under desiccating conditions. The product can be stored for up to 12 months.
溶解性	Soluble in DMSO to 100 mM and in ethanol to 100 mM
使用に関する注意	<p>Wherever possible, you should prepare and use solutions on the same day. However, if you need to make up stock solutions in advance, we recommend that you store the solution as aliquots in tightly sealed vials at -20°C. Generally, these will be useable for up to one month. Before use, and prior to opening the vial we recommend that you allow your product to equilibrate to room temperature for at least 1 hour.</p> <p>Toxic, refer to SDS for further information.</p> <p>Need more advice on solubility, usage and handling? Please visit our frequently asked questions</p>

[\(FAQ\) page](#) for more details.

SMILES 線形表記

Clc1cc(NN=C(/C#N)C#N)ccc1

由来

Synthetic

アプリケーション


The Abpromise guarantee

Abpromise保証は、次のテスト済みアプリケーションにおけるab141229の使用に適用されます

アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。


アプリケーション	Abreviews	特記事項
Functional Studies		Use at an assay dependent concentration.

画像



2D chemical structure image of ab141229, CCCP, Mitochondrial oxidative phosphorylation uncoupler

Chemical Structure - CCCP, Mitochondrial oxidative phosphorylation uncoupler (ab141229)




Immunocytochemistry/ Immunofluorescence -
CCCP, Mitochondrial oxidative phosphorylation
uncoupler (ab141229)

Immunofluorescent analysis of 4 % paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized HeLa (human epithelial cell line from cervix adenocarcinoma)(+/- treatment with 10µM carbonyl cyanide 3-chlorophenylhydrazine (CCCP, ab141229) for 24 hours) cells labeling PINK1 with **ab216144** at 1/500 dilution, followed by Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) (**ab150077**) secondary antibody at 1/1000 dilution (green). Confocal image showing cytoplasmic staining on HeLa cells treated with 10µM carbonyl cyanide 3-chlorophenylhydrazine (CCCP, ab141229) for 24 hours. The nuclear counter stain is DAPI (blue). Tubulin is detected with Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) (**ab195889**) at 1/200 dilution (red).

The negative controls are as follows:

-ve control: PBS, followed by Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) (**ab150077**) secondary antibody at 1/1000 dilution.



Western blot - CCCP, Mitochondrial oxidative
phosphorylation uncoupler (ab141229)

All lanes : Anti-PINK1 antibody [EPR20730] (**ab216144**) at 1/1000 dilution

Lane 1 : HeLa (human epithelial cell line from cervix adenocarcinoma) whole cell lysate

Lane 2 : HeLa cells (treated with 10uM carbonyl cyanide 3-chlorophenylhydrazine (CCCP, ab141229) for 24 hours) whole cell lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/20000 dilution


Developed using the ECL technique.

Observed band size: 62 kDa

Exposure time: 5 seconds

Blocking and dilution buffer: 5% NFDm/TBST

PINK1 can be induced by CCCP treatment (PMID: 24184327).



Immunoprecipitation - CCCP, Mitochondrial oxidative phosphorylation uncoupler (ab141229)


PINK1 was immunoprecipitated from 0.35 mg of HeLa (human epithelial cell line from cervix adenocarcinoma) (treated with 10uM carbonyl cyanide 3-chlorophenylhydrazone (CCCP. ab141229) for 24 hours) whole cell lysate with **ab216144** at 1/30 dilution. Western blot was performed from the immunoprecipitate using **ab216144** at 1/500 dilution. VeriBlot for IP Detection Reagent (HRP) (**ab131366**), was used for detection at 1/1000 dilution.

Lane 1: HeLa (CCCP-treated, ab141229) lysate 10 µg (Input).

Lane 2: **ab216144** IP in HeLa (CCCP-treated, ab141229) lysate.

Lane 3: Rabbit monoclonal IgG (**ab172730**) instead of **ab216144** in HeLa (CCCP-treated, ab141229) whole cell lysate.

Blocking and dilution buffer: 5% NFDm/TBST.



Functional Studies - CCCP, Mitochondrial oxidative phosphorylation uncoupler (ab141229)

MCF7 cells were incubated at 37°C for 2 hours with vehicle control (0 µM) and different concentrations of CCCP (ab 141229). Increased expression of AKT1 (phospho S473) (**ab81283**) in MCF7 cells correlates with an increase in CCCP concentration, as described in literature.

Whole cell lysates were prepared with RIPA buffer (containing protease inhibitors and sodium orthovanadate), 10 µg of each were loaded on the gel and the WB was run under reducing conditions. After transfer the membrane was blocked for an hour using 5% BSA before being incubated with **ab81283** at 2 µg/ml and **ab8227** at 1 µg/ml overnight at 4°C. Antibody binding was detected using an anti-rabbit antibody conjugated to HRP (**ab97051**) at 1/10000 and visualised using ECL development solution.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES, NOT FOR USE IN HUMANS"

Our Promise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Promise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors
- Abcam biochemicals are novel compounds and we have not tested their biological activity in house. Please use the literature to identify how to use these products effectively. If you require further assistance please contact the scientific support team