

Exo-1, Vesicular trafficking inhibitor ab120292


[2 References](#) [画像数 2](#)

製品の概要

製品名	Exo-1, Vesicular trafficking inhibitor
製品の詳細	Vesicular trafficking inhibitor
生理活性の詳細	Cell-permeable, reversible inhibitor of vesicular trafficking between endoplasmic reticulum and the Golgi apparatus. Inhibits exocytosis with an IC ₅₀ of approx. 20 μM. Rapidly releases Arf1 from Golgi membranes. Has similar mechanism of action to brefeldin A (ab120299).

CAS 番号 461681-88-9

構造式




製品の特性

体系名	Methyl 2-(4-fluorobenzamido)benzoate
分子量	273.26
分子式	C ₁₅ H ₁₂ FNO ₃
PubChem 登録番号	310557
保存方法	Store at Room Temperature. The product can be stored for up to 12 months.
溶解性	Soluble in DMSO to 100 mM and in ethanol to 25 mM
使用に関する注意	Wherever possible, you should prepare and use solutions on the same day. However, if you need to make up stock solutions in advance, we recommend that you store the solution as aliquots in tightly sealed vials at -20°C. Generally, these will be useable for up to one month. Before use, and prior to opening the vial we recommend that you allow your product to equilibrate to room temperature for at least 1 hour.

Need more advice on solubility, usage and handling? Please visit our [frequently asked questions \(FAQ\) page](#) for more details.


SMILES 線形表記 O=C(Nc1cccc1C(=O)OC)c2ccc(F)cc2

由来 Synthetic



Chemical Structure - Exo-1, Vesicular trafficking inhibitor (ab120292)

2D chemical structure image of ab120292, Exo-1, Vesicular trafficking inhibitor



Immunocytochemistry/ Immunofluorescence - Exo-1, Vesicular trafficking inhibitor (ab120292)

ab86071 staining GBF1 in HeLa cells treated with Exo-1 (ab120292), by ICC/IF. Increase in GBF1 expression correlates with increased concentration of Exo-1 as described in literature. The cells were incubated at 37°C for 5 minutes in media containing different concentrations of ab120292 (Exo-1) in DMSO, fixed with 4% formaldehyde for 10 minutes at room temperature and blocked with PBS containing 10% goat serum, 0.3 M glycine, 1% BSA and 0.1% tween for 2h at room temperature. Staining of the treated cells with **ab86071** (5 μg/ml) was performed overnight at 4°C in PBS containing 1% BSA and 0.1% tween. A DyLight 488 goat anti-rabbit polyclonal antibody (**ab96899**) at 1/250 dilution was used as the secondary antibody. Nuclei were counterstained with DAPI and are shown in blue.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES, NOT FOR USE IN HUMANS"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise,

please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors
- Abcam biochemicals are novel compounds and we have not tested their biological activity in house. Please use the literature to identify how to use these products effectively. If you require further assistance please contact the scientific support team