


XAV939, Tankyrase inhibitor ab120897

8 References [画像数 2](#)

製品の概要

製品名	XAV939, Tankyrase inhibitor
製品の詳細	Potent tankyrase inhibitor. Inhibitor of Wnt/ β -catenin signaling.
精製度	> 98%
CAS 番号	284028-89-3
構造式	



製品の特性

体系名	3,5,7,8-Tetrahydro-2-[4-(trifluoromethyl)phenyl]-4 <i>H</i> -thiopyrano[4,3- <i>d</i>]pyrimidin-4-one
分子量	312.31
分子式	C ₁₄ H ₁₁ F ₃ N ₂ OS
保存方法	Store at Room Temperature. The product can be stored for up to 12 months.
溶解性	Soluble in DMSO to 25 mM
使用に関する注意	<p>Wherever possible, you should prepare and use solutions on the same day. However, if you need to make up stock solutions in advance, we recommend that you store the solution as aliquots in tightly sealed vials at -20°C. Generally, these will be useable for up to one month. Before use, and prior to opening the vial we recommend that you allow your product to equilibrate to room temperature for at least 1 hour.</p> <p>Toxic, refer to SDS for further information.</p> <p>Need more advice on solubility, usage and handling? Please visit our frequently asked questions (FAQ) page for more details.</p>
SMILES 線形表記	OC1=NC(C2=CC=C(C(F)(F)F)C=C2)=NC3=C1CSCC3
由来	Synthetic

アプリケーション


The Abpromise guarantee

Abpromise保証は、次のテスト済みアプリケーションにおけるab120897の使用に適用されます

アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。


アプリケーション	Abreviews	特記事項
Functional Studies		Use at an assay dependent concentration.

画像



2D chemical structure image of ab120897, XAV939, Tankyrase inhibitor

Chemical Structure - XAV939, Tankyrase inhibitor (ab120897)



Immunocytochemistry/ Immunofluorescence - XAV939, Tankyrase inhibitor (ab120897)

ab16051 staining β -catenin in SW480 cells treated with XAV939 (ab120897), by ICC/IF. Increase of β -catenin cytoplasmic expression and decrease in nuclear expression correlates with increased concentration of XAV939, as described in literature. The cells were incubated at 37°C for 6 hours in media containing different concentrations of ab120897 (XAV939) in DMSO, fixed with 4% formaldehyde for 10 minutes at room temperature and blocked with PBS containing 10% goat serum, 0.3 M glycine, 1% BSA and 0.1% tween for 2h at room temperature. Staining of the treated cells with **ab16051** (1 μ g/ml) was performed overnight at 4°C in PBS containing 1% BSA and 0.1% tween. A DyLight 488 anti-rabbit polyclonal antibody (**ab96899**) at 1/250 dilution was used as the secondary antibody. Nuclei were counterstained with DAPI and are shown in blue.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES, NOT FOR USE IN HUMANS"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors
- Abcam biochemicals are novel compounds and we have not tested their biological activity in house. Please use the literature to identify how to use these products effectively. If you require further assistance please contact the scientific support team