

### Jurkat whole cell lysate ab7899

#### 3 References

#### 製品の概要

製品名	Jurkat whole cell lysate
特記事項	<p>Cell line: Jurkat, clone E6-1 (Human acute T cell leukemia).</p> <p>Growth media: RPMI and 10% NCS (Newborn calf serum).</p> <p>Jurkat cell lysate was prepared by homogenization in modified RIPA buffer (150 mM sodiumchloride, 50 mM Tris-HCl, pH 7.4, 1 mM ethylenediaminetetraacetic acid, 1 mM phenylmethylsulfonyl flouride, 1% Triton X-100, 1% sodium deoxycholic acid, 0.1% sodium dodecylsulfate, 5 µg/ml of aprotinin, 5 µg/ml of leupeptin). Cell debris was removed by centrifugation. Protein concentration was determined with Bio-Rad protein assay. The cell lysate was boiled for 5 min in 1 x SDS sample buffer (0.045 M Tris-HCl pH 6.8, 10% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue), containing 0.05 M DTT.</p>
アプリケーション	適用あり: WB

#### 製品の特性

Mycoplasma free	Yes
製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot. Store at -80°C. Avoid freeze / thaw cycle.
バッファー	<p>pH: 7.20</p> <p>Constituent: 100% SDS Sample Buffer</p>
ライゼート 備考	<p>Jurkat cell lysate was prepared by homogenization in modified RIPA buffer (150 mM sodiumchloride, 50 mM Tris-HCl, pH 7.4, 1 mM ethylenediaminetetraacetic acid, 1 mM phenylmethylsulfonyl flouride, 1% Triton X-100, 1% sodium deoxycholic acid, 0.1% sodium dodecylsulfate, 5 µg/ml of aprotinin, 5 µg/ml of leupeptin). Cell debris was removed by centrifugation. Protein concentration was determined with Bio-Rad protein assay. The cell lysate was boiled for 5 min in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol.</p>
背景	<p>Often this cell line is called 'JM' (Jurkat and JM are derived from the same patient and are sister clones), occasionally JM may be a subclone with somewhat divergent features. Jurkat cells can be transfected and are therefore useful for studies of blood proto-oncogenes expression, apoptosis and cell survival (e.g HIV). They produce IL-2 and can also be used for studies of differentiation.</p>

## アプリケーション

---

**The Abpromise guarantee**      Abpromise保証は、 次のテスト済みアプリケーションにおけるab7899の使用に適用されます  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		Use at an assay dependent dilution. Jurkat cell lysate is <b>ready to load</b> on SDS-PAGE for Western blotting, 20 µg per lane is recommended for mini gel.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

## Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors