# abcam

## Product datasheet

# MOLT-4 whole cell lysate ab7912

#### 製品の概要

製品名 MOLT-4 whole cell lysate

特記事項 Cell line: MOLT4 (Human acute lymphoblastic leukemia).

Growth media: RPMI and 10% FBS.

MOLT4 cell lysate was prepared by homogenization in modified RIPA buffer (150 mM sodium chloride, 50 mM Tris-HCl, pH 7.4, 1 mM ethylenediaminetetraacetic acid, 1 mM phenylmethylsulfonyl flouride, 1% Triton X-100, 1% sodium deoxycholic acid, 0.1% sodium dodecylsulfate, 5  $\mu$ g/ml of aprotinin, 5  $\mu$ g/ml of leupeptin). Cell debris was removed by centrifugation. Protein concentration was determined with Bio-Rad protein assay. The cell lysate was boiled for 5 min in 1 x SDS sample buffer (0.045 M Tris-HCl pH 6.8, 10% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue), containing 0.05 M DTT.

アプリケーション **適用あり**: WB

#### 製品の特性

Mycoplasma free Yes

製品の状態 Liquid

保存方法 Shipped at 4°C. Upon delivery aliquot. Store at -80°C. Avoid freeze / thaw cycle.

**バッファー** pH: 7.20

Constituent: 100% SDS Sample Buffer

ライゼート 備考 MOLT4 cell lysate was prepared by homogenization in modified RIPA buffer (150 mM sodium

chloride, 50 mM Tris-HCl, pH 7.4, 1 mM ethylenediaminetetraacetic acid, 1 mM

 $phenylmethyl sulfonyl \ flouride, \ 1\% \ Triton \ X-100, \ 1\% \ sodium \ deoxycholic \ acid, \ 0.1\% \ sodium$ 

dodecylsulfate,  $5 \mu g/ml$  of aprotinin,  $5 \mu g/ml$  of leupeptin). Cell debris was removed by centrifugation. Protein concentration was determined with Bio-Rad protein assay. The cell lysate

was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1%  $\,$ 

sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol.

背景 MOLT4 cells are lymphoblast-like in morphology and are used for studies of apoptosis, tumor

cytotoxicity, tumorigenicity, as well as for antitumour testing.

アプリケーション

1

### The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab7912の使用に適用されます

アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		Use at an assay dependent concentration.  MOLT4 cell lysate is ready to load on SDS-PAGE for Western blotting, 20 µg per lane is recommended for mini gel.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

#### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- · Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="https://www.abcam.co.jp/abpromise">https://www.abcam.co.jp/abpromise</a> or contact our technical team.

#### Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors