abcam

Product datasheet

Human Perforin ELISPOT Set ab46582

製品の概要

製品の概要

製品名 Human Perforin ELISPOT Set

サンプルの種類 Suspension cells

アッセイタイプ Sandwich (qualitative)

ステップ Multiple steps standard assay

種交差性 交差種: Human

AZE: Haman

The Human Perforin ELISPOT Set is designed to enumerate Perforin producing cells in a single cell suspension. This method has the advantage of requiring a minimum of in-vitro manipulations allowing perforin production analysis as close as possible to in-vivo conditions in a highly specific way. This technique is designed to determine the frequency of perforin producing cells under a given stimulation, and the follow-up of such frequency during a treatment and/or a pathological state. Elispot assay constitutes an ideal tool in the TH1 / TH2 response, vaccine development, viral infection monitoring and treatment, cancerology, infectious diseases, autoimmune diseases and transplantation.

This ELISPOT assay is based on sandwich immuno-enzyme technology. Cell secreted cytokines or soluble molecules are captured by coated antibodies avoiding diffusion in supernatant, protease degradation or binding on soluble membrane receptors. After cell removal, the captured cytokines are revealed by tracer antibodies and appropriate conjugates.

Principal:

After cell stimulation, locally produced cytokines are captured by a specific monoclonal antibody. After cell lysis, trapped cytokine molecules are revealed by a secondary biotinylated detection antibody, which is in turn recognised by streptavidin conjugated to alkaline phosphatase. PVDF-bottomed-well plates are then incubated with BCIP/NBT substrate. Colored "purple" spots indicate cytokine production by individual cells.

Recognizes natural human Perforin.

アプリケーション 適用あり: ELISpot

試験プラットフォーム Reagents

製品の特性

1

保存方法

Store at +4°C. Please refer to protocols.

内容	5 x 96 tests	10 x 96 tests	15 x 96 tests	20 x 96 tests
Bovine Serum Albumin	1 x 1g	2 x 2g	3 x 1g	4 x 1g
Human Perforin Capture Antibody	1 x 500µl	2 x 500µl	3 x 500µl	4 x 500µl
Perforin Biotinylated Detection Antibody	1 vial	2 vials	3 vials	4 vials
Ready-to-use BCIP/NBT substrate buffer	1 x 50ml	4 x 27ml	3 x 50ml	4 x 50ml
Streptavidin - Alkaline Phosphatase conjugated	1 x 50µl	2 x 50µl	3 x 50µl	4 x 50μl

関連性

Perforin has structural and functional similarities to complement component 9 (C9). Like C9, this protein creates transmembrane tubules and is capable of lysing non-specifically a variety of target cells. Perforin is one of the main cytolytic proteins of cytolytic granules, and it is known to be a key effector molecule for T-cell- and natural killer-cell-mediated cytolysis. Defects in this gene cause familial hemophagocytic lymphohistiocytosis type 2 (HPLH2), a rare and lethal autosomal recessive disorder of early childhood.

細胞内局在

Cell Membrane and Cytoplasmic

アプリケーション

The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab46582の使用に適用されますアプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
ELISpot		Use at an assay dependent dilution.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

•	Guarantee only valid for products bought direct from Abcam or one of our authorized distributors						
		3					