

### Anti-Adiponectin antibody ab62551

★★★★★ **1 Abreviews**   **15 References**   画像数 **2**

#### 製品の概要

|              |   |
|--------------|---|
| 製品名          | Anti-Adiponectin antibody   |
| 製品の詳細        | Rabbit polyclonal to Adiponectin  |
| 由来種          | Rabbit  |
| アプリケーション     | <b>適用あり:</b> WB, IHC-P  |
| 種交差性         | <b>交差種:</b> Rat   |
| 免疫原          | A 15 amino acid peptide from near the amino terminus of human Adiponectin.  |
| ポジティブ・コントロール | Rat brain cell lysate. Rat brain tissue.  |
| 特記事項         | <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p> |

#### 製品の特性

|        |   |
|--------|---|
| 製品の状態  | Liquid  |
| 保存方法   | Shipped at 4°C. Store at +4°C.                                  |
| バッファー  | pH: 7.2<br>Preservative: 0.02% Sodium azide<br>Constituent: PBS |
| 精製度    | Immunogen affinity purified                                     |
| ポリ/モノ  | ポリクローナル   |
| アイソタイプ | IgG   |

#### アプリケーション


**The Abpromise guarantee**   **Abpromise保証は、** 次のテスト済みアプリケーションにおけるab62551の使用に適用されます  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

| アプリケーション | Abreviews | 特記事項   |
|----------|-----------|--|
| WB       | ★★★★★ (1) | Use a concentration of 1 µg/ml. Detects a band of approximately 32 kDa (predicted molecular weight: 26 kDa). |
| IHC-P    |           | Use a concentration of 1 µg/ml.  |

## ターゲット情報

|       |   |
|-------|---|
| 機能    | Important adipokine involved in the control of fat metabolism and insulin sensitivity, with direct anti-diabetic, anti-atherogenic and anti-inflammatory activities. Stimulates AMPK phosphorylation and activation in the liver and the skeletal muscle, enhancing glucose utilization and fatty-acid combustion. Antagonizes TNF-alpha by negatively regulating its expression in various tissues such as liver and macrophages, and also by counteracting its effects. Inhibits endothelial NF-kappa-B signaling through a cAMP-dependent pathway. May play a role in cell growth, angiogenesis and tissue remodeling by binding and sequestering various growth factors with distinct binding affinities, depending on the type of complex, LMW, MMW or HMW.  |
| 組織特異性 | Synthesized exclusively by adipocytes and secreted into plasma.   |
| 関連疾患  | Defects in ADIPOQ are the cause of adiponectin deficiency (ADPND) [MIM:612556]. ADPND results in very low concentrations of plasma adiponectin.<br>Genetic variations in ADIPOQ are associated with non-insulin-dependent diabetes mellitus (NIDDM) [MIM:125853]; also known as diabetes mellitus type 2. NIDDM is characterized by an autosomal dominant mode of inheritance, onset during adulthood and insulin resistance.   |
| 配列類似性 | Contains 1 C1q domain.<br>Contains 1 collagen-like domain.  |
| ドメイン  | The C1q domain is commonly called the globular domain.  |
| 翻訳後修飾 | Hydroxylated Lys-33 was not identified in PubMed:16497731, probably due to poor representation of the N-terminal peptide in mass fingerprinting.<br>HMW complexes are more extensively glycosylated than smaller oligomers. Hydroxylation and glycosylation of the lysine residues within the collagen-like domain of adiponectin seem to be critically involved in regulating the formation and/or secretion of HMW complexes and consequently contribute to the insulin-sensitizing activity of adiponectin in hepatocytes.<br>O-glycosylated. Not N-glycosylated. O-linked glycans on hydroxylysines consist of Glc-Gal disaccharides bound to the oxygen atom of post-translationally added hydroxyl groups. Sialylated to varying degrees depending on tissue. Thr-22 appears to be the major site of sialylation. Higher sialylation found in SGBS adipocytes than in HEK fibroblasts. Sialylation is not required neither for heterodimerization nor for secretion. Not sialylated on the glycosylated hydroxylysines.<br>Desialylated forms are rapidly cleared from the circulation. |
| 細胞内局在 | Secreted.   |

## 画像



**Lane 1 :** Anti-Adiponectin antibody (ab62551) at 1 µg/ml

**Lane 2 :** Anti-Adiponectin antibody (ab62551) at 2 µg/ml


**Lane 3 :** Anti-Adiponectin antibody (ab62551) at 4 µg/ml

**All lanes :** Rat brain cell lysate

Lysates/proteins at 15 µg per lane.

**Predicted band size:** 26 kDa

**Observed band size:** 32 kDa



ab62551, at 1µg/ml, staining Adiponectin in rat brain tissue by immunohistochemistry using paraffin-embedded tissue.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

## Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors