

Anti-Agmatine antibody ab62667

医薬用外劇物

製品の概要

製品名	Anti-Agmatine antibody
製品の詳細	Chicken polyclonal to Agmatine
由来種	Chicken
特異性	Fixed tissue cross-reactivity tested with known targets at recommended dilution. No measurable glutaraldehyde-fixed tissue cross-reactivity (<1:1000) against L-alanine, gamma-aminobutyrate, L-citrulline, L-cysteine, D/L-glutamate, D/L-arginine, D/L-glutamine, glutathione, glycine, L-lysine, L-ornithine, L-serine, taurine, L-threonine, L-tryptophan, L-tyrosine.
アプリケーション	適用あり: ICC
種交差性	交差種: Species independent
免疫原	Agmatine cross-linked to purified fraction V bovine serum albumin with glutaraldehyde.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	Preservative: 0.05% Thimerosal (merthiolate) Constituent: 0.268% PBS
精製度	IgY fraction
特記事項(精製)	IgYs were separated from other yolk proteins by serial PEG6000 extractions using a modification of the method of Polsonet al Immunological Investigations. 1985, 14, 323-327.
ポリモノ	ポリクローナル
アイソタイプ	IgY

アプリケーション

The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab62667の使用に適用されます**
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
ICC		

追加情報

ICC: 1/100. Excitation mapping by LM immunocytochemistry. True dilution at user dilution: 1/2000. Optimal fixation: 0.1-2.5% glutaraldehyde, 1% formaldehyde using HPI (High Performance Immunocytochemistry). Minimum glutaraldehyde: 0.1% using HPI.

Not yet tested in other applications.

Optimal dilutions/concentrations should be determined by the end user.

ターゲット情報

関連性

Agmatine, a product of arginine metabolism is an endogenous amine acting at imidazoline receptors with a putative role in nociception at the spinal level. Agmatine potentiates morphine analgesia and blocks symptoms of naloxone-precipitated morphine withdrawal in rats.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors