

Anti-ALDH1A1 antibody [EP1933Y] - BSA and Azide free ab215996

リコンビナント **RabMAb**

★★★★★ **1 Abreviews** **25 References** 画像数 **14**

製品の概要

製品名	Anti-ALDH1A1 antibody [EP1933Y] - BSA and Azide free
製品の詳細	Rabbit monoclonal [EP1933Y] to ALDH1A1 - BSA and Azide free
由来種	Rabbit
特異性	The mouse recommendation is based on the WB results. We do not guarantee IHC-P for mouse.
アプリケーション	適用あり: Flow Cyt (Intra), ICC/IF, WB, IP, IHC-P
種交差性	交差種: Mouse, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	IP: HepG2 cell lysate; IHC-P: Human liver tissue, Human bladder carcinomaFlow Cyt (intra): HepG2 cells
特記事項	ab215996 is the carrier-free version of <u>ab52492</u> .

Our **carrier-free** antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.

This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our **conjugation kits** for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

This product is compatible with the Maxpar[®] Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar[®] is a trademark of Fluidigm Canada Inc.

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information **see here**.

Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit

monoclonal antibodies. For details on our patents, please refer to [RabMAb® patents](#).

Rat: We have preliminary internal testing data to indicate this antibody may not react with this species. Please contact us for more information.

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C. Do Not Freeze.
バッファー	pH: 7.20 Constituent: PBS
キャリア・フリー	はい
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	EP1933Y
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee **Abpromise保証は、**次のテスト済みアプリケーションにおけるab215996の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
Flow Cyt (Intra)		Use at an assay dependent concentration. ab199376 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.
ICC/IF		Use at an assay dependent concentration.
WB		Use at an assay dependent concentration. Detects a band of approximately 55 kDa (predicted molecular weight: 55 kDa).
IP		Use at an assay dependent concentration.
IHC-P	★☆☆☆☆ (1)	Use at an assay dependent concentration. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol. See IHC antigen retrieval protocols . The mouse recommendation is based on the WB results. We do not guarantee IHC-P for mouse.

ターゲット情報

機能	Binds free retinal and cellular retinol-binding protein-bound retinal. Can convert/oxidize retinaldehyde to retinoic acid.
----	--

パスウェイ

Cofactor metabolism; retinol metabolism.


配列類似性

Belongs to the aldehyde dehydrogenase family.

細胞内局在


Cytoplasm.

画像



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human breast carcinoma tissue sections labeling ALDH1A1 with purified [ab52492](#) at 1/50 dilution (3.54 µg/ml). Perform heat mediated antigen retrieval using [ab93684](#) (Tris/EDTA buffer, pH 9.0). ImmunoHistoProbe one step HRP Polymer (ready to use) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab52492](#)).



[ab52492](#) (purified) at 1/20 dilution (2ug) immunoprecipitating ALDH1A1 in HepG2 whole cell lysates.

Lane 1: HepG2 (Human hepatocellular carcinoma epithelial cell) whole cell lysates 10ug


Lane 2 (+): [ab52492](#) & HepG2 whole cell lysates

Lane 3 (-): Rabbit monoclonal IgG ([ab172730](#)) instead of [ab52492](#) in HepG2 whole cell lysates

For western blotting, VeriBlot for IP Detection Reagent (HRP) ([ab131366](#)) was used at 1/1000 dilution.

Blocking and diluting buffer: 5% NFDM/TBST.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab52492](#)).



Immunocytochemistry/ Immunofluorescence - Anti-ALDH1A1 antibody [EP1933Y] - BSA and Azide free (ab215996)


Immunocytochemistry/ Immunofluorescence analysis of HepG2 (Human liver hepatocellular carcinoma cell line) cells labeling ALDH1A1 with [ab52492](#) (purified) at 1/500 dilution (4 µg/ml).

Cells were fixed in 100% methanol. [ab150077](#), an AlexaFluor®488 Goat anti-Rabbit secondary antibody was used at 1/1000 dilution (2 µg/ml). [ab195889](#), Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) was used to counterstain at 1/200 dilution (2.5 µg/ml). DAPI was used as nuclear counterstain.

Confocal image showing cytoplasmic staining on HepG2 cell line.

Negative control: No staining on MCF-7 cell line.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab52492](#)).




Flow Cytometry (Intracellular) - Anti-ALDH1A1 antibody [EP1933Y] - BSA and Azide free (ab215996)

Clone EP1933Y (ab215996) has been successfully conjugated by Abcam. This image was generated using Anti-ALDH1A1 antibody [EP1933Y] (PE). Please refer to [ab209437](#) for protocol details.

Overlay histogram showing MCF7 cells stained with [ab209437](#) (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Triton X-100 for 15 min at 22°C. The cells were then incubated in 1x PBS / 10% normal goat serum to block non-specific protein-protein interactions followed by the antibody ([ab209437](#), 1/500 dilution) for 30 min at 22°C.

Isotype control antibody (black line) was rabbit IgG (monoclonal) Phycoerythrin ([ab209478](#)) used at the same concentration and conditions as the primary antibody. Unlabelled sample (blue line) was also used as a control.

Acquisition of >5,000 events were collected using a 50 mW Yellow/Green laser (561nm) and 586/15 bandpass filter.




Immunocytochemistry/ Immunofluorescence - Anti-ALDH1A1 antibody [EP1933Y] - BSA and Azide free (ab215996)

Clone EP1933Y (ab215996) has been successfully conjugated by Abcam. This image was generated using Anti-ALDH1A1 antibody [EP1933Y] (Alexa Fluor® 647). Please refer to [ab195255](#) for protocol details.

ab195255 staining ALDH1A1 in MCF7 cells. The cells were fixed with 4% formaldehyde (10 min), permeabilized in 0.1% Triton X-100 for 5 minutes and then blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated with **ab195255** at a working dilution of 1 in 50 (shown in red) and **ab195887**, Mouse monoclonal [DM1A] to alpha Tubulin (Alexa Fluor® 488, shown in green) at 2µg/ml overnight at +4°C. Nuclear DNA was labelled in blue with DAPI.

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).




Immunocytochemistry/ Immunofluorescence - Anti-ALDH1A1 antibody [EP1933Y] - BSA and Azide free (ab215996)

Clone EP1933Y (ab215996) has been successfully conjugated by Abcam. This image was generated using Anti-ALDH1A1 antibody [EP1933Y] (Alexa Fluor® 488). Please refer to [ab195254](#) for protocol details.

ab195254 staining ALDH1A1 in MCF7 cells. The cells were fixed with 4% formaldehyde (10 min), permeabilised in 0.1% Triton X-100 for 5 minutes and then blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated with **ab195254** at a working dilution of 1 in 100 (shown in green) and **ab195889**, Mouse monoclonal [DM1A] to alpha Tubulin (Alexa Fluor® 594, shown in red) at 2µg/ml overnight at +4°C. Nuclear DNA was labelled in blue with DAPI.


Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).



Flow Cytometry (Intracellular) - Anti-ALDH1A1 antibody [EP1933Y] - BSA and Azide free (ab215996)

Intracellular Flow Cytometry analysis of HepG2 (Human hepatocellular carcinoma epithelial cell) cells labeling ALDH1A1 with purified [ab52492](#) at 1/20 dilution (10µg/ml) (red). Cells were fixed with 4% Paraformaldehyde and permeabilised with 90% Methanol. A Goat anti rabbit IgG (Alexa Fluor® 488, [ab150077](#)) secondary antibody was used at 1/2000. Isotype control - Rabbit monoclonal IgG (Black). Unlabeled control - Cell without incubation with primary antibody and secondary antibody (Blue).

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab52492](#)).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ALDH1A1 antibody [EP1933Y] - BSA and Azide free (ab215996)

Image from Gong et al PLoS One. 2010 Dec 20;5(12):e15630. doi: 10.1371/journal.pone.0015630. Fig 1.


Tumor tissues of primary invasive ductal carcinomas of the breast were obtained from 192 female patients with stage IIB and III prior to pre-operative neoadjuvant chemotherapy.

(progressive or stable disease, PD/SD)

(partial or complete remission, PR/CR)

The level of ALDH1 was tested by immunohistochemistry staining in paraffin-embedded tissue sections. Rabbit monoclonal ALDH1A1 antibody ([ab52492](#), unpurified, Abcam) used at a 1:100 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab52492](#)).




Flow Cytometry (Intracellular) - Anti-ALDH1A1 antibody [EP1933Y] - BSA and Azide free (ab215996)

Overlay histogram showing HepG2 (Human liver hepatocellular carcinoma cell line) cells stained with **ab52492** (unpurified) (red line).

The cells were fixed with 80% methanol (5 minutes) and then permeabilized with 0.1% PBS-Tween for 20 minutes. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (**ab52492**, 1/1000 dilution) for 30 minutes at 22°C. The secondary antibody used was Alexa Fluor® 488 goat anti-rabbit IgG (H&L) (**ab150077**) at 1/2000 dilution for 30 minutes at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (0.1 µg/1x10⁶ cells) used under the same conditions. Unlabeled sample (blue line) was also used as a control.

Acquisition of >5,000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter.


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab52492**).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ALDH1A1 antibody [EP1933Y] - BSA and Azide free (ab215996)

Immunohistochemical analysis of paraffin-embedded human liver tissue sections labeling ALDH1A1 with **ab52492** (unpurified) at 1/100 dilution.


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab52492**).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ALDH1A1 antibody [EP1933Y] - BSA and Azide free (ab215996)

Immunohistochemical analysis of Formalin/PFA-fixed paraffin-embedded normal human brain tissue sections labeling ALDH1A1 with **ab52492** (unpurified).


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab52492**).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ALDH1A1 antibody [EP1933Y] - BSA and Azide free (ab215996)

Immunohistochemical analysis of Formalin/PFA-fixed paraffin-embedded normal human lung tissue sections labeling ALDH1A1 with **ab52492** (unpurified).

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab52492**).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ALDH1A1 antibody [EP1933Y] - BSA and Azide free (ab215996)

Immunohistochemical analysis of Formalin/PFA-fixed paraffin-embedded normal human kidney tissue sections labeling ALDH1A1 with **ab52492** (unpurified).

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab52492**).

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-ALDH1A1 antibody [EP1933Y] - BSA and Azide free (ab215996)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors