abcam

Product datasheet

Alexa Fluor® 488 Anti-PABPN1 antibody [EP3000Y] ab206056

ועלשעבע RabMAb

画像数 2 1 References

製品の概要

製品名 Alexa Fluor® 488 Anti-PABPN1 antibody [EP3000Y]

製品の詳細 Alexa Fluor® 488 Rabbit monoclonal [EP3000Y] to PABPN1

由来種 Rabbit

標識 Alexa Fluor® 488. Ex: 495nm, Em: 519nm

アプリケーション 適用あり: ICC/IF 種交差性 交差種: Human

交差が予測される動物種: Mouse 4

免疫原 Synthetic peptide within Human PABPN1 (N terminal). The exact sequence is proprietary.

Database link: Q86U42

ポジティブ・コントロール ICC/IF: HeLa cells

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit 特記事項

monoclonal antibodies. For details on our patents, please refer to **RabMAb**® patents. Alexa Fluor[®] is a registered trademark of Molecular Probes, Inc., a Thermo Fisher Scientific

Company. The Alexa Fluor[®] dye included in this product is provided under an intellectual property license from Life Technologies Corporation. As this product contains the Alexa Fluor® dye, the purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). As this product contains the Alexa Fluor[®] dye the sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: in manufacturing; (ii) to provide a service, information, or data in return for payment (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are sold for use in research. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, 5781 Van Allen Way, Carlsbad, CA 92008 USA or

outlicensing@thermofisher.com.

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long

term. Avoid freeze / thaw cycle. Store In the Dark.

バッファー pH: 7.40

Preservative: 0.02% Sodium azide

Constituents: PBS, 30% Glycerol (glycerin, glycerine), 1% BSA

精製度 Affinity purified ポリ/モノ モノクローナル PDローン名 EP3000Y

アイソタイプ IgG

アプリケーション

The Abpromise guarantee <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab206056の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
ICC/IF		1/100. This product gave a positive signal in HeLa cells fixed with 4% formaldehyde (10 min).

ターゲット情報

機能 Involved in the 3'-end formation of mRNA precursors (pre-mRNA) by the addition of a poly(A) tail

of 200-250 nt to the upstream cleavage product. Stimulates poly(A) polymerase (PAPOLA) conferring processivity on the poly(A) tail elongation reaction and controls also the poly(A) tail length. Increases the affinity of poly(A) polymerase for RNA. Is also present at various stages of mRNA metabolism including nucleocytoplasmic trafficking and nonsense-mediated decay (NMD) of mRNA. Cooperates with SKIP to synergistically activate E-box-mediated transcription through MYOD1 and may regulate the expression of muscle-specific genes. Binds to poly(A) and to

poly(G) with high affinity. May protect the poly(A) tail from degradation.

組織特異性 Ubiquitous.

関連疾患 Defects in PABPN1 are the cause of oculopharyngeal muscular dystrophy (OPMD) [MIM:164300].

OPMD is a form of late-onset slowly progressive myopathy characterized by eyelid ptosis,

dysphagia and, sometimes by other cranial and limb-muscle involvement.

配列類似性 Contains 1 RRM (RNA recognition motif) domain.

ドメイン The RRM domain is essential for specific adenine bases recognition in the poly(A) tail but not

sufficient for poly(A) binding.

翻訳後修飾 Arginine dimethylation is asymmetric and involves PRMT1 and PRMT3. It does not influence the

RNA binding properties.

細胞内局在 Nucleus. Cytoplasm. Localized in cytoplasmic mRNP granules containing untranslated mRNAs.


Shuttles between the nucleus and the cytoplasm but predominantly found in the nucleus. Its nuclear import may involve the nucleocytoplasmic transport receptor transportin and a RAN-GTP-sensitive

import mechanism. Is exported to the cytoplasm by a carrier-mediated pathway that is

independent of mRNA traffic. Nucleus; nuclear speckle. Colocalizes with SKIP and poly(A) RNA in

nuclear speckles.


画像



Immunocytochemistry/ Immunofluorescence - Alexa Fluor® 488 Anti-PABPN1 antibody [EP3000Y] (ab206056)

ab206056 staining PABPN1 in HeLa cells. The cells were fixed with 4% formaldehyde (10 min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at +4°C with ab206056 at 1/100 dilution (shown in green) and ab195889, Mouse monoclonal to alpha Tubulin (Alexa Fluor[®] 594), at 1/250 dilution (shown in red). Nuclear DNA was labelled with DAPI (shown in blue).

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- · Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish

- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors