abcam

Product datasheet

Alexa Fluor® 555 Anti-IKB alpha antibody [E130] ab206578

יעלאעבע RabMAb

画像数 2

製品の概要

製品名 Alexa Fluor® 555 Anti-IKB alpha antibody [E130]

製品の詳細 Alexa Fluor® 555 Rabbit monoclonal [E130] to IKB alpha

Rabbit

Alexa Fluor® 555. Ex: 555nm, Em: 565nm

適用あり: ICC/IF 交差種: Human

交差が予測される動物種: Mouse, Rat, Hamster, Cow, Pig

Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

ICC/IF: HeLa cells

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

Alexa Fluor[®] is a registered trademark of Molecular Probes, Inc., a Thermo Fisher Scientific Company. The Alexa Fluor® dye included in this product is provided under an intellectual property license from Life Technologies Corporation. As this product contains the Alexa Fluor® dye, the purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). As this product contains the Alexa Fluor[®] dye the sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: in manufacturing; (ii) to provide a service, information, or data in return for payment (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are sold for use in research. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, 5781 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@thermofisher.com.

由来種

標識

アプリケーション

種交差性

免疫原

ポジティブ・コントロール

特記事項

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C.

Avoid freeze / thaw cycle. Store In the Dark.

バッファー pH: 7.40

Preservative: 0.02% Sodium azide

Constituents: 30% Glycerol (glycerin, glycerine), 1% BSA, PBS

精製度 Protein A purified

ポリ/モノ モノクローナル

クローン名 E130 **アイソタイプ** IgG

アプリケーション

The Abpromise guarantee <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab206578の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
ICC/IF		1/500. This product gave a positive signal in HeLa cells fixed with 100% methanol (5 min)

ターゲット情報

機能 Inhibits the activity of dimeric NF-kappa-B/REL complexes by trapping REL dimers in the

cytoplasm through masking of their nuclear localization signals. On cellular stimulation by immune

and proinflammatory responses, becomes phosphorylated promoting ubiquitination and

degradation, enabling the dimeric RELA to translocate to the nucleus and activate transcription.

関連疾患 Ectodermal dysplasia, anhidrotic, with T-cell immunodeficiency autosomal dominant

配列類似性 Belongs to the NF-kappa-B inhibitor family.

Contains 5 ANK repeats.

翻訳後修飾 Phosphorylated; disables inhibition of NF-kappa-B DNA-binding activity. Phosphorylation at

positions 32 and 36 is prerequisite to recognition by UBE2D3 leading to polyubiquitination and

subsequent degradation.

Sumoylated; sumoylation requires the presence of the nuclear import signal. Sumoylation blocks

 $ubiquitination\ and\ proteasome-mediated\ degradation\ of\ the\ protein\ thereby\ increasing\ the\ protein$

stability.

Monoubiquitinated at Lys-21 and/or Lys-22 by UBE2D3. Ubiquitin chain elongation is then performed by CDC34 in cooperation with the SCF(FBXW11) E3 ligase complex, building

ubiquitin chains from the UBE2D3-primed NFKBIA-linked ubiquitin. The resulting


polyubiquitination leads to protein degradation. Also ubiquitinated by SCF(BTRC) following

stimulus-dependent phosphorylation at Ser-32 and Ser-36.

Deubiquitinated by porcine reproductive and respiratory syndrome virus Nsp2 protein, which

Cytoplasm. Nucleus. Shuttles between the nucleus and the cytoplasm by a nuclear localization signal (NLS) and a CRM1-dependent nuclear export.

画像



Immunocytochemistry/ Immunofluorescence - Alexa Fluor® 555 Anti-IKB alpha antibody [E130] (ab206578) ab206578 staining IKB alpha in HeLa cells. The cells were fixed with 100% methanol (5min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at +4°C with ab206578 at 1/500 dilution (pseudocolored in green) and ab195884, Rat monoclonal to Tubulin (Alexa Fluor® 647), at 1/250 dilution (shown in red). Nuclear DNA was labelled with DAPI (shown in blue).

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).



 $\textbf{Please note:} \ \ \textbf{All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"}$

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery

- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors