

Alexa Fluor® 647 Anti-CD69 antibody [FN50] ab187772

画像数 1

製品の概要

| | |
|--------------|---|
| 製品名 | Alexa Fluor® 647 Anti-CD69 antibody [FN50] |
| 製品の詳細 | Alexa Fluor® 647 Mouse monoclonal [FN50] to CD69 |
| 由来種 | Mouse |
| 標識 | Alexa Fluor® 647. Ex: 652nm, Em: 668nm |
| アプリケーション | 適用あり: Flow Cyt |
| 種交差性 | 交差種: Human |
| 免疫原 | Tissue, cells or virus corresponding to Human CD69. anti-μ-stimulated human B lymphocytes |
| ポジティブ・コントロール | Human blood cells |
| 特記事項 | <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p> |

製品の特性

| | |
|--------|--|
| 製品の状態 | Liquid |
| 保存方法 | Shipped at 4°C. Store at +4°C. Do Not Freeze. Store In the Dark. |
| バッファー | <p>pH: 7.4</p> <p>Preservative: 0.097% Sodium azide</p> <p>Constituents: 0.2% BSA, 99% PBS</p> |
| 精製度 | Size exclusion |
| ポリ/モノ | モノクローナル |
| クローン名 | FN50 |
| アイソタイプ | IgG1 |

アプリケーション

The Abpromise guarantee

Abpromise保証は、次のテスト済みアプリケーションにおけるab187772の使用に適用されます


アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

| アプリケーション | Abreviews | 特記事項 |
|----------|-----------|--|
| Flow Cyt | | Use 4µl for 10 ⁶ cells. or 100 µl of whole blood |

ターゲット情報

| | |
|-------|---|
| 機能 | Involved in lymphocyte proliferation and functions as a signal transmitting receptor in lymphocytes, natural killer (NK) cells, and platelets. |
| 組織特異性 | Expressed on the surface of activated T-cells, B-cells, natural killer cells, neutrophils, eosinophils, epidermal Langerhans cells and platelets. |
| 配列類似性 | Contains 1 C-type lectin domain. |
| 発生段階 | Earliest inducible cell surface glycoprotein acquired during lymphoid activation. |
| 翻訳後修飾 | Constitutive Ser/Thr phosphorylation in both mature thymocytes and activated T-lymphocytes. |
| 細胞内局在 | Membrane. |

画像



Flow Cytometry analysis of Human peripheral blood cells labeling CD69 with Anti-CD69 antibody [FN50] (Alexa Fluor® 647) (ab187772).

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors