abcam

Product datasheet

Alexa Fluor® 647 Anti-IKB alpha antibody [EP697] ab207356

יעלאעבע RabMAb

画像数 2

製品の概要

製品名 Alexa Fluor® 647 Anti-IKB alpha antibody [EP697]

製品の詳細 Alexa Fluor® 647 Rabbit monoclonal [EP697] to IKB alpha

由来種 Rabbit

標識 Alexa Fluor® 647. Ex: 652nm, Em: 668nm

アプリケーション 適用あり: ICC/IF 種交差性 交差種: Human

交差が予測される動物種: Mouse, Rat 4

Synthetic peptide within Human IKB alpha aa 300 to the C-terminus. The exact immunogen sequence used to generate this antibody is proprietary information. If additional detail on the immunogen is needed to determine the suitability of the antibody for your needs, please contact

our Scientific Support team to discuss your requirements.

Database link: P25963

Run BLAST with Run BLAST with

ポジティブ・コントロール

特記事項

免疫原

ICC/IF: HeLa cells

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

Alexa Fluor[®] is a registered trademark of Molecular Probes, Inc, a Thermo Fisher Scientific Company. The Alexa Fluor[®] dye included in this product is provided under an intellectual property license from Life Technologies Corporation. As this product contains the Alexa Fluor® dye, the purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). As this product contains the Alexa Fluor[®] dye the sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: in manufacturing; (ii) to provide a service, information, or data in return for payment (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are sold for use in research. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, 5781 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@thermofisher.com.

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C.

Stable for 12 months at -20°C. Store In the Dark.

バッファー pH: 7.40

Preservative: 0.02% Sodium azide

Constituents: 30% Glycerol (glycerin, glycerine), 1% BSA, PBS

精製度 Protein A purified

ポリ/モノ モノクローナル

ウローン名 EP697 **アイソタイプ** IgG

アプリケーション

The Abpromise guarantee <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab207356の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
ICC/IF		1/50. This product gave a positive signal in HeLa cells fixed with 4% formaldehyde (10 min)

ターゲット情報

機能 Inhibits the activity of dimeric NF-kappa-B/REL complexes by trapping REL dimers in the

cytoplasm through masking of their nuclear localization signals. On cellular stimulation by immune

and proinflammatory responses, becomes phosphorylated promoting ubiquitination and

degradation, enabling the dimeric RELA to translocate to the nucleus and activate transcription.

関連疾患 Ectodermal dysplasia, anhidrotic, with T-cell immunodeficiency autosomal dominant

配列類似性 Belongs to the NF-kappa-B inhibitor family.

Contains 5 ANK repeats.

翻訳後修飾 Phosphorylated; disables inhibition of NF-kappa-B DNA-binding activity. Phosphorylation at

positions 32 and 36 is prerequisite to recognition by UBE2D3 leading to polyubiquitination and

subsequent degradation.

Sumoylated; sumoylation requires the presence of the nuclear import signal. Sumoylation blocks

ubiquitination and proteasome-mediated degradation of the protein thereby increasing the protein

stability

Monoubiquitinated at Lys-21 and/or Lys-22 by UBE2D3. Ubiquitin chain elongation is then performed by CDC34 in cooperation with the SCF(FBXW11) E3 ligase complex, building


ubiquitin chains from the UBE2D3-primed NFKBIA-linked ubiquitin. The resulting

polyubiquitination leads to protein degradation. Also ubiquitinated by SCF(BTRC) following


stimulus-dependent phosphorylation at Ser-32 and Ser-36.

Deubiquitinated by porcine reproductive and respiratory syndrome virus Nsp2 protein, which thereby interferes with NFKBIA degradation and impairs subsequent NF-kappa-B activation.

画像



Immunocytochemistry/ Immunofluorescence - Alexa Fluor® 647 Anti-IKB alpha antibody [EP697] (ab207356) ab207356 staining IKB alpha in HeLa cells. The cells were fixed with 4% formaldehyde (10 min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at +4°C with ab207356 at 1/50 dilution (shown in red) and ab195887, Mouse monoclonal to alpha Tubulin (Alexa Fluor® 488), at 1/250 dilution (shown in green). Nuclear DNA was labelled with DAPI (shown in blue). Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors