

Anti-AP2 gamma/TFAP2C antibody [3B5] ab110635

1 Abreviews 1 References 画像数 5

製品の概要

製品名	Anti-AP2 gamma/TFAP2C antibody [3B5]
製品の詳細	Mouse monoclonal [3B5] to AP2 gamma/TFAP2C
由来種	Mouse
アプリケーション	適用あり: IHC-P, WB, ICC/IF
種交差性	交差種: Mouse, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	WB: PC-7, A431, mouse placenta, and RAW 264.7 cell lysates; ICC/IF: HeLa, MCF-7, and U251 cells; IHC-P: Human breast cancer;
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
バッファー	Preservative: 0.05% Sodium azide Constituents: 99.85% PBS, 0.1% BSA
精製度	Protein G purified
ポリ/モノ	モノクローナル
クローン名	3B5
アイソタイプ	IgG2b

アプリケーション

The Abpromise guarantee Abpromise保証は、 次のテスト済みアプリケーションにおけるab110635の使用に適用されます


アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
IHC-P		1/50.
WB		1/100 - 1/1000. Predicted molecular weight: 49 kDa. Block with BSA only (not milk).
ICC/IF		1/50 - 1/500.

ターゲット情報


機能	Sequence-specific DNA-binding protein that interacts with inducible viral and cellular enhancer elements to regulate transcription of selected genes. AP-2 factors bind to the consensus sequence 5'-GCCNNNGGC-3' and activate genes involved in a large spectrum of important biological functions including proper eye, face, body wall, limb and neural tube development. They also suppress a number of genes including MCAM/MUC18, C/EBP alpha and MYC.
配列類似性	Belongs to the AP-2 family.
ドメイン	The WW-binding motif mediates interaction with WWOX.
翻訳後修飾	Sumoylated on Lys-10; which inhibits transcriptional activity.
細胞内局在	Nucleus.

画像



Immunofluorescent analysis of AP2 gamma/TFAP2C using ab110635 shows staining in HeLa Cells. AP2 gamma/TFAP2C staining (green), F-Actin staining with Phalloidin (red) and nuclei with DAPI (blue) is shown. Cells were grown on chamber slides and fixed with formaldehyde prior to staining. Cells were probed without (control) or with ab110635 at a dilution of 1:100 over night at 4°C, washed with PBS and incubated with a Goat Anti-Mouse DyLight-488 conjugated secondary antibody. Images were taken at 60X magnification.

Immunocytochemistry/ Immunofluorescence - Anti-AP2 gamma/TFAP2C antibody [3B5] (ab110635)



Western blot - Anti-AP2 gamma/TFAP2C antibody [3B5] (ab110635)

All lanes : Anti-AP2 gamma/TFAP2C antibody [3B5] (ab110635) at 1/50 dilution

Lane 1 : PC-3 cell lysate

Lane 2 : A431 cell lysate

Lane 3 : Mouse placenta cell lysate


Lane 4 : RAW 264.7 cell lysate

Lysates/proteins at 30 µg per lane.

Secondary


All lanes : Goat anti-Mouse IgG (H+L), HRP conjugate at 1/4000 dilution

Predicted band size: 49 kDa




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-AP2 gamma/TFAP2C antibody [3B5] (ab110635)

Immunohistochemistry was performed on cancer biopsies of deparaffinized Human breast carcinoma tissue. To expose target proteins, heat induced antigen retrieval was performed using 10mM sodium citrate (pH6.0) buffer, microwaved for 8-15 minutes. Following antigen retrieval tissues were blocked in 3% BSA-PBS for 30 minutes at room temperature. Tissues were then probed at a dilution of 1:50 with ab110635 or without primary antibody (negative control) overnight at 4°C in a humidified chamber. Tissues were washed extensively with PBST and endogenous peroxidase activity was quenched with a peroxidase suppressor. Detection was performed using a biotin-conjugated secondary antibody and SA-HRP, followed by colorimetric detection using DAB. Tissues were counterstained with hematoxylin and prepped for mounting.



Immunocytochemistry/ Immunofluorescence - Anti-AP2 gamma/TFAP2C antibody [3B5] (ab110635)

Immunofluorescent analysis of AP2 gamma/TFAP2C using ab110635 shows staining in U251 Cells. AP2 gamma/TFAP2C staining (green), F-Actin staining with Phalloidin (red) and nuclei with DAPI (blue) is shown. Cells were grown on chamber slides and fixed with formaldehyde prior to staining. Cells were probed without (control) or with ab110635 at a dilution of 1:100 over night at 4°C, washed with PBS and incubated with a Goat Anti-Mouse DyLight-488 conjugated secondary antibody. Images were taken at 60X magnification.



Immunocytochemistry/ Immunofluorescence - Anti-AP2 gamma/TFAP2C antibody [3B5] (ab110635)

Immunofluorescent analysis of AP2 gamma/TFAP2C using ab110635 shows staining in MCF-7 Cells. AP2 gamma/TFAP2C staining (green), F-Actin staining with Phalloidin (red) and nuclei with DAPI (blue) is shown. Cells were grown on chamber slides and fixed with formaldehyde prior to staining. Cells were probed without (control) or with ab110635 at a dilution of 1:20 over night at 4°C, washed with PBS and incubated with a Goat Anti-Mouse DyLight-488 conjugated secondary antibody. Images were taken at 60X magnification.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors