

APC Anti-CCR7 antibody [4B12] ab93518

画像数 1

製品の概要

製品名	APC Anti-CCR7 antibody [4B12]
製品の詳細	APC Rat monoclonal [4B12] to CCR7
由来種	Rat
標識	APC. Ex: 645nm, Em: 660nm
アプリケーション	適用あり: Flow Cyt
種交差性	交差種: Mouse
免疫原	Tissue, cells or virus corresponding to CCR7. Rat basophilic leukaemia (RBL)-2H3 cells transfected with Mouse CCR7.
ポジティブ・コントロール	C57Bl/6 splenocytes
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C.
バッファー	pH: 7.20 Preservative: 0.09% Sodium azide Constituents: 0.87% Sodium chloride, PBS
精製度	Affinity purified
ポリ/モノ	モノクローナル
クローン名	4B12
アイソタイプ	IgG2a
軽鎖の種類	kappa

アプリケーション


The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab93518の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
Flow Cyt		Use 1 µg for 10 ⁶ cells.

ターゲット情報

機能	Receptor for the MIP-3-beta chemokine. Probable mediator of EBV effects on B-lymphocytes or of normal lymphocyte functions.
組織特異性	Expressed in various lymphoid tissues and activated B- and T-lymphocytes, strongly up-regulated in B-cells infected with Epstein-Barr virus and T-cells infected with herpesvirus 6 or 7.
配列類似性	Belongs to the G-protein coupled receptor 1 family.
細胞内局在	Cell membrane.

画像



Staining of C57Bl/6 splenocytes with FITC anti-mouse CD4 (GK1.5) and either 0.5 µg of ab93518 (right) or 0.5 µg of APC Rat IgG2a Iso Control (left). Cells were stained at 37°C for 0.5 hours and Fc receptor binding was blocked by staining cells with 5 µg/million cells anti-mouse CD16/32. Cells in the lymphocyte gate were used for analysis.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors