

### APC Anti-CD11c antibody [N418] ab210306

画像数 1

#### 製品の概要

製品名	APC Anti-CD11c antibody [N418]
製品の詳細	APC Armenian hamster monoclonal [N418] to CD11c
由来種	Armenian hamster
標識	APC. Ex: 645nm, Em: 660nm
アプリケーション	<b>適用あり:</b> Flow Cyt
種交差性	<b>交差種:</b> Mouse
免疫原	Tissue, cells or virus corresponding to Mouse CD11c.
ポジティブ・コントロール	C57Bl/6 splenocytes.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

#### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C. Store In the Dark.
バッファー	<p>pH: 7.20</p> <p>Preservative: 0.09% Sodium azide</p> <p>Constituents: 0.12% Monobasic dihydrogen sodium phosphate, 0.87% Sodium chloride, 0.1% Gelatin</p>
精製度	Affinity purified
特記事項(精製)	ab210306 was purified from tissue culture supernatant via affinity chromatography. The purified antibody was conjugated under optimal conditions, with unreacted dye removed from the preparation.
ポリ/モノ	モノクローナル
クローン名	N418

## アプリケーション

### The Abpromise guarantee

**Abpromise保証は、**次のテスト済みアプリケーションにおけるab210306の使用に適用されます

アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
Flow Cyt		Use at an assay dependent concentration. Use 0.125 µg.

## ターゲット情報

### 機能

Integrin alpha-X/beta-2 is a receptor for fibrinogen. It recognizes the sequence G-P-R in fibrinogen. It mediates cell-cell interaction during inflammatory responses. It is especially important in monocyte adhesion and chemotaxis.

### 組織特異性

Predominantly expressed in monocytes and granulocytes.

### 配列類似性

Belongs to the integrin alpha chain family.  
Contains 7 FG-GAP repeats.  
Contains 1 VWFA domain.


### ドメイン

The integrin I-domain (insert) is a VWFA domain. Integrins with I-domains do not undergo protease cleavage.

### 細胞内局在

Membrane.

## 画像



Flow cytometric analysis of C57Bl/6 splenocytes stained with an Anti-Mouse MHC Class II FITC, followed by labeling of CD11c with 0.125 µg ab210306 (right panel), compared with 0.125 µg Armenian Hamster IgG Allophycocyanin (left panel).

### **Our Abpromise to you: Quality guaranteed and expert technical support**

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors