

# APC Anti-CD69 antibody [H1.2F3] ab95611

### 画像数 1

#### 製品の概要

---

製品名	APC Anti-CD69 antibody [H1.2F3]
製品の詳細	APC Armenian hamster monoclonal [H1.2F3] to CD69
由来種	Armenian hamster
標識	APC. Ex: 645nm, Em: 660nm
アプリケーション	<b>適用あり:</b> Flow Cyt
種交差性	<b>交差種:</b> Mouse
免疫原	Tissue, cells or virus corresponding to Mouse CD69. Murine dendritic epidermal T cell line Y245.
ポジティブ・コントロール	1-day ConA-stimulated BALB/c splenocytes; stimulated Mouse splenic cell cultures
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

#### 製品の特性

---

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C.
バッファー	pH: 7.20 Preservative: 0.09% Sodium azide
精製度	Protein G purified
ポリ/モノ	モノクローナル
クローン名	H1.2F3
アイソタイプ	IgG

#### アプリケーション

---


**The Abpromise guarantee** **Abpromise保証は、次のテスト済みアプリケーションにおけるab95611の使用に適用されます**  
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
Flow Cyt		Use 0.125-0.25µg for 10 <sup>5-8</sup> cells. Staining the cell sample in a final volume of 100 µl is recommended.

## ターゲット情報

<b>機能</b>	Involved in lymphocyte proliferation and functions as a signal transmitting receptor in lymphocytes, natural killer (NK) cells, and platelets.
<b>組織特異性</b>	Expressed on the surface of activated T-cells, B-cells, natural killer cells, neutrophils, eosinophils, epidermal Langerhans cells and platelets.
<b>配列類似性</b>	Contains 1 C-type lectin domain.
<b>発生段階</b>	Earliest inducible cell surface glycoprotein acquired during lymphoid activation.
<b>翻訳後修飾</b>	Constitutive Ser/Thr phosphorylation in both mature thymocytes and activated T-lymphocytes.
<b>細胞内局在</b>	Membrane.

## 画像



Staining of 1-day ConA-stimulated BALB/c splenocytes with 0.125 µg of Armenian Hamster IgG Isotype Control APC (blue histogram) or 0.125 µg of ab95611 (purple histogram). Total viable cells were used for analysis.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

## Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors