abcam

Product datasheet

Anti-ATP5I antibody ab122241

4 References 画像数 3

製品の概要

製品名 Anti-ATP5I antibody

製品の詳細 Rabbit polyclonal to ATP5I

由来種 Rabbit

アプリケーション 適用あり: ICC/IF, IHC-P, WB

種交差性 交差種: Human 免疫原 antigen sequence:

RRIAAEEKKKQDELKRIARELAEDDSILK

, corresponding to amino acids 41-69 of Human ATP5I.

Run BLAST with
Run BLAST with

ポジティブ・コントロール Human stomach tissue; RT-4 and U-251 MG cell lysates; Human Liver and Tonsil lysates; Human

cell line A-431

特記事項

ab122241 is mono-specific.

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

バッファー pH: 7.20

Preservative: 0.02% Sodium azide Constituents: 40% Glycerol, 59% PBS

精製度 Immunogen affinity purified 一次抗体 備考 ab122241 is mono-specific.

1

アイソタイプ

ΙgG

アプリケーション

The Abpromise guarantee <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab122241の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
ICC/IF		Use a concentration of 0.25 - 2 µg/ml.
IHC-P		1/200 - 1/500. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
WB		Use a concentration of 0.04 - 0.4 μg/ml.

ターゲット情報

機能

Mitochondrial membrane ATP synthase (F(1)F(0) ATP synthase or Complex V) produces ATP from ADP in the presence of a proton gradient across the membrane which is generated by electron transport complexes of the respiratory chain. F-type ATPases consist of two structural domains, F(1) - containing the extramembraneous catalytic core, and F(0) - containing the membrane proton channel, linked together by a central stalk and a peripheral stalk. During catalysis, ATP synthesis in the catalytic domain of F(1) is coupled via a rotary mechanism of the central stalk subunits to proton translocation. Part of the complex F(0) domain. Minor subunit located with subunit a in the membrane.

配列類似性

Belongs to the ATPase e subunit family.

細胞内局在

Mitochondrion. Mitochondrion inner membrane.

画像



All lanes: Anti-ATP5I antibody (ab122241) at 1/250 dilution


Lane 1: RT-4 cell lysate

Lane 2: U-251 MG cell lysate

Lane 3: Human Plasma lysate


Lane 4: Human Liver lysate

Lane 5: Human Tonsil lysate



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-ATP5I antibody (ab122241)

ab122241, at 1/200 dilution, staining ATP5I in paraffin-embedded Human stomach tissue by Immunohistochemistry.



Immunocytochemistry/ Immunofluorescence - Anti-ATP5I antibody (ab122241)

ab122241, at 4µg/ml, staining ATP5I in Human cell line A-431 by Immunofluorescence. Antibody staining is shown in green.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors