

### Anti-BarX2 antibody [BAR X2 8A7/1] ab20128

★★★★☆ [1 Abreviews](#) [1 References](#)

#### 製品の概要

製品名	Anti-BarX2 antibody [BAR X2 8A7/1]
製品の詳細	Mouse monoclonal [BAR X2 8A7/1] to BarX2
由来種	Mouse
アプリケーション	<b>適用あり:</b> ELISA, IHC-P, ICC/IF, WB
種交差性	<b>交差種:</b> Human
免疫原	Fusion protein: MBP-tagged BarX2 (Human).
特記事項	

BarX2 is a tumour suppressor gene that encodes for a transcription factor known to regulate transcription of cell adhesion molecules in the mouse.

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

#### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	Preservative: 0.02% Sodium azide Constituent: 99.98% PBS
精製度	Protein A/G purified
一次抗体 備考	BarX2 is a tumour suppressor gene that encodes for a transcription factor known to regulate transcription of cell adhesion molecules in the mouse.
ポリ/モノ	モノクローナル
クローン名	BAR X2 8A7/1
ミエローマ	Sp2/0-Ag14

アイソタイプ	IgG1
軽鎖の種類	unknown

## アプリケーション

**The Abpromise guarantee** **Abpromise保証は、次のテスト済みアプリケーションにおけるab20128の使用に適用されず**  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
ELISA		Use at an assay dependent concentration.
IHC-P		Use at an assay dependent concentration.
ICC/IF		Use at an assay dependent concentration.
WB	★★★★★ (1)	Use at an assay dependent concentration.

## ターゲット情報

機能	Transcription factor. Binds optimally to the DNA consensus sequence 5'-YYTAATGRTTTTY-3'. May control the expression of neural adhesion molecules such as L1 or Ng-CAM during embryonic development of both the central and peripheral nervous system. May be involved in controlling adhesive processes in keratinizing epithelia.
組織特異性	Highly expressed in adult salivary gland and at much lower levels in mammary gland, kidney and placenta.
配列類似性	Belongs to the BAR homeobox family. Contains 1 homeobox DNA-binding domain.
細胞内局在	Nucleus.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

## Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors