

Anti-Basophils antibody [2D7] ab155577

★★★★★ [6 Abreviews](#) [6 References](#)

製品の概要

製品名	Anti-Basophils antibody [2D7]
製品の詳細	Mouse monoclonal [2D7] to Basophils
由来種	Mouse
アプリケーション	適用あり: WB, IHC-P, Flow Cyt, IHC-Fr, ICC/IF
種交差性	交差種: Human
免疫原	Tissue, cells or virus corresponding to Human Basophils.
ポジティブ・コントロール	Chronic myeloid leukaemia (basophilia). Negative control: CD203c negative cells (Basophil depleted cell fractions).
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	Preservative: 0.02% Sodium azide Constituents: 99% PBS, 0.1% BSA
精製度	Protein G purified
ポリ/モノ	モノクローナル
クローン名	2D7
アイソタイプ	IgG1

アプリケーション

The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab155577の使用に適用されます
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
WB	★★★★★ (1)	Use at an assay dependent concentration.
IHC-P	★★★★★ (1)	Use at an assay dependent concentration. Fresh surgical tissues can be fixed in Carnoy's fluid for 24 h and transferred to absolute ethanol.
Flow Cyt	★★★★★ (1)	Use at an assay dependent concentration. ab170190 - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody.
IHC-Fr	★★★★★ (2)	Use at an assay dependent concentration. Fix in either 10% NBF or Carnoy's fluid for 15 minutes at RT.
ICC/IF	★★★★★ (1)	Use at an assay dependent concentration. Cytospin preparations should be fixed in methanol for 15 min, incubated in methanol containing 0.6% H ₂ O ₂ , for 30 min to block endogenous peroxidase.

ターゲット情報

関連性

Basophils are multifunctional haematopoietic cells that secrete a wide variety of proinflammatory agents upon activation such as histamine and cytokines. Moreover, basophils can induce IgE production by B cells. Under normal conditions basophils are primarily produced in the bone marrow, whereas mature basophils reside in the circulation and can enter tissues at sites of inflammation. Both basophils and mast cells have been implicated in the pathogenesis of allergic inflammation due to the abundant expression of high affinity receptors for IgE. The 2D7 antibody can be used as a sensitive and precise marker for human basophils and does not react with lymphocytes, monocytes, eosinophils, neutrophils or mast cells in immunohistochemistry. Moreover, the antibody is suitable for western blot analysis of the 2D7 antigen.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors