


Anti-Bax antibody ab104156

KO 評価済

[12 References](#) [画像数 3](#)

製品の概要

製品名	Anti-Bax antibody
製品の詳細	Rabbit polyclonal to Bax
由来種	Rabbit
特異性	Expression levels of BAX protein vary with sample type. Induction may be required if endogenous expression is low.
アプリケーション	適用あり: WB
種交差性	交差種: Human 交差が予測される動物種: Cow, Cat, Dog, Pig, Macaque monkey 
免疫原	Synthetic peptide within Human Bax aa 1-100 conjugated to keyhole limpet haemocyanin. The exact sequence is proprietary. Database link: Q07812
ポジティブ・コントロール	This antibody gave a positive signal in Human lymph node tissue lysate.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	pH: 7.40 Preservative: 0.02% Sodium azide Constituent: PBS
	Batches of this product that have a concentration < 1mg/ml may have BSA added as a stabilising

agent. If you would like information about the formulation of a specific lot, please contact our scientific support team who will be happy to help.

精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション


The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab104156の使用に適用されず
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		Use a concentration of 1 µg/ml. Detects a band of approximately 21 kDa (predicted molecular weight: 21 kDa).

ターゲット情報

機能	Accelerates programmed cell death by binding to, and antagonizing the apoptosis repressor BCL2 or its adenovirus homolog E1B 19k protein. Under stress conditions, undergoes a conformation change that causes translocation to the mitochondrion membrane, leading to the release of cytochrome c that then triggers apoptosis. Promotes activation of CASP3, and thereby apoptosis.
組織特異性	Expressed in a wide variety of tissues. Isoform Psi is found in glial tumors. Isoform Alpha is expressed in spleen, breast, ovary, testis, colon and brain, and at low levels in skin and lung. Isoform Sigma is expressed in spleen, breast, ovary, testis, lung, colon, brain and at low levels in skin. Isoform Alpha and isoform Sigma are expressed in pro-myelocytic leukemia, histiocytic lymphoma, Burkitt's lymphoma, T-cell lymphoma, lymphoblastic leukemia, breast adenocarcinoma, ovary adenocarcinoma, prostate carcinoma, prostate adenocarcinoma, lung carcinoma, epidermoid carcinoma, small cell lung carcinoma and colon adenocarcinoma cell lines.
配列類似性	Belongs to the Bcl-2 family.
ドメイン	Intact BH3 motif is required by BIK, BID, BAK, BAD and BAX for their pro-apoptotic activity and for their interaction with anti-apoptotic members of the Bcl-2 family.
細胞内局在	Cytoplasm and Mitochondrion membrane. Cytoplasm. Colocalizes with 14-3-3 proteins in the cytoplasm. Under stress conditions, undergoes a conformation change that causes release from JNK-phosphorylated 14-3-3 proteins and translocation to the mitochondrion membrane.

画像



Western blot - Anti-Bax antibody (ab104156)

Lanes 1, 3 and 5: Wild-type HAP1 cell lysate (20 µg)


Lanes 2, 4 and 6: Bax knockout HAP1 cell lysate (20 µg)

Lanes 1 and 2: Green signal from target – ab104156 observed at 20 kDa

Lanes 3 and 4: Red signal from loading control – **ab8226** observed at 42 kDa

Lanes 5 and 6: Merged (red and green) signal

ab104156 was shown to react with Bax when Bax knockout samples were used, along with additional cross-reactive bands. Wild-type and Bax knockout samples were subjected to SDS-PAGE. ab104156 and **ab8226** (loading control to beta actin) were diluted 1 µg/mL and 1/2000 respectively and incubated overnight at 4°C. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed **ab216773** and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed **ab216776** secondary antibodies at 1/10000 dilution for 1 h at room temperature before imaging.



Western blot - Anti-Bax antibody (ab104156)

Anti-Bax antibody (ab104156) at 1 µg/ml + Human lymph node tissue lysate - total protein (**ab29871**) at 10 µg

Secondary

Goat Anti-Rabbit IgG H&L (HRP) preadsorbed (**ab97080**) at 1/5000 dilution


Developed using the ECL technique.

Performed under reducing conditions.

Predicted band size: 21 kDa

Observed band size: 21 kDa

Exposure time: 4 minutes



Lane 1: Wild-type HAP1 cell lysate (20 µg)

Lane 2: Bax knockout HAP1 cell lysate (20 µg)

Lanes 1 - 2: Merged signal (red and green). Green - ab104156 observed at 20 kDa. Red - loading control, **ab8245**, observed at 37 kDa or **ab18058**, observed at 130 kDa.

This western blot image is a comparison between ab104156 and a competitor's top cited rabbit polyclonal antibody.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors