




### Anti-Bcl-2 antibody [100/D5] ab692

KO 評価済

★★★★★ 9 Abreviews 243 References 画像数 4

#### 製品の概要

製品名	Anti-Bcl-2 antibody [100/D5]
製品の詳細	Mouse monoclonal [100/D5] to Bcl-2
由来種	Mouse
アプリケーション	適用あり: WB, Flow Cyt, IHC-P, ICC/IF
種交差性	交差種: Human 交差が予測される動物種: Cow, Dog, Chinese hamster  非交差種: Rat
免疫原	Synthetic peptide corresponding to Bcl-2 aa 41-54. Sequence: GAAPAPGIFSSQPG-Cys  Database link: <b>P10415</b> <div>  <a href="#">Run BLAST with</a>  <a href="#">Run BLAST with</a> </div>
ポジティブ・コントロール	IHC: Human tonsil ICC/IF: Human neuroblastoma (SK-N-SH cells) WB: HAP1 cells and HeLa cells Flow Cyt: Jurkat cells
特記事項	<p>This product was changed from ascites to tissue culture supernatant on 8<sup>th</sup> March 2018. Please note that the dilutions may need to be adjusted accordingly. If you have any questions, please do not hesitate to contact our scientific support team.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

#### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle.

バッファー	pH: 7.30 Preservative: 0.09% Sodium azide Constituents: PBS, Tissue culture supernatant, 1% BSA  Proprietary preservative that is not sodium azide or thimerosal, protein carrier.
精製度	Tissue culture supernatant
ポリ/モノ	モノクローナル
クローン名	100/D5
ミエローマ	P3-NS1/1-Ag4-1
アイソタイプ	IgG1
軽鎖の種類	kappa

## アプリケーション

**The Abpromise guarantee**      **Abpromise保証は、次のテスト済みアプリケーションにおけるab692の使用に適用されます**  
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB	★★★★★ (7)	Use at an assay dependent concentration. Predicted molecular weight: 26 kDa.
Flow Cyt		1/10. <b>ab170190</b> - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody.
IHC-P		Use at an assay dependent concentration.
ICC/IF		Use a concentration of 5 µg/ml.

## ターゲット情報

機能	Suppresses apoptosis in a variety of cell systems including factor-dependent lymphohematopoietic and neural cells. Regulates cell death by controlling the mitochondrial membrane permeability. Appears to function in a feedback loop system with caspases. Inhibits caspase activity either by preventing the release of cytochrome c from the mitochondria and/or by binding to the apoptosis-activating factor (APAF-1). May attenuate inflammation by impairing NLRP1-inflammasome activation, hence CASP1 activation and IL1B release (PubMed:17418785).
組織特異性	Expressed in a variety of tissues.
関連疾患	A chromosomal aberration involving BCL2 has been found in chronic lymphatic leukemia. Translocation t(14;18)(q32;q21) with immunoglobulin gene regions. BCL2 mutations found in non-Hodgkin lymphomas carrying the chromosomal translocation could be attributed to the Ig somatic hypermutation mechanism resulting in nucleotide transitions.
配列類似性	Belongs to the Bcl-2 family.
ドメイン	BH1 and BH2 domains are required for the interaction with BAX and for anti-apoptotic activity.

The BH4 motif is required for anti-apoptotic activity and for interaction with RAF1 and EGLN3. The loop between motifs BH4 and BH3 is required for the interaction with NLRP1.


## 翻訳後修飾

Phosphorylation/dephosphorylation on Ser-70 regulates anti-apoptotic activity. Growth factor-stimulated phosphorylation on Ser-70 by PKC is required for the anti-apoptosis activity and occurs during the G2/M phase of the cell cycle. In the absence of growth factors, BCL2 appears to be phosphorylated by other protein kinases such as ERKs and stress-activated kinases. Phosphorylated by MAPK8/JNK1 at Thr-69, Ser-70 and Ser-87, which stimulates starvation-induced autophagy. Dephosphorylated by protein phosphatase 2A (PP2A). Proteolytically cleaved by caspases during apoptosis. The cleaved protein, lacking the BH4 motif, has pro-apoptotic activity, causes the release of cytochrome c into the cytosol promoting further caspase activity. Monoubiquitinated by PARK2, leading to increase its stability. Ubiquitinated by SCF(FBXO10), leading to its degradation by the proteasome.

## 細胞内局在

Mitochondrion outer membrane. Nucleus membrane. Endoplasmic reticulum membrane.

## 画像




**Lane 1:** Wild type HAP1 whole cell lysate (20 µg)

**Lane 2:** BCL2 knockout HAP1 whole cell lysate (20 µg)

**Lane 3:** HeLa whole cell lysate (20 µg)

**Lanes 1 - 3:** Merged signal (red and green). Green - ab692 observed at 26 kDa. Red - loading control, [ab181602](#), observed at 37 kDa.

ab692 was shown to specifically react with BCL2 when BCL2 knockout samples were used. Wild-type and BCL2 knockout samples were subjected to SDS-PAGE. Ab692 and [ab181602](#) (Rabbit anti GAPDH loading control) were incubated overnight at 4°C at 500 dilution and 1/10000 dilution respectively. Blots were developed with Goat anti-Mouse IgG H&L (IRDye® 800CW) preabsorbed [ab216772](#) and Goat anti-Rabbit IgG H&L (IRDye® 680RD) preabsorbed [ab216777](#) secondary antibodies at 1/10000 dilution for 1 hour at room temperature before imaging.




ab692 staining Bcl-2 in SK-N-SH cells treated with (R)-(-)-Deprenyl hydrochloride (Selegiline hydrochloride) ([ab120604](#)), by ICC/IF.

Increase of Bcl-2 expression correlates with increased concentration of (R)-(-)-Deprenyl hydrochloride (Selegiline hydrochloride), as described in literature.


The cells were incubated at 37°C for 3h in media containing different concentrations of [ab120604](#) ((R)-(-)-Deprenyl hydrochloride (Selegiline hydrochloride)) in DMSO, fixed with 4% formaldehyde for 10 minutes at room temperature and blocked with PBS containing 10% goat serum, 0.3 M glycine, 1% BSA and 0.1%

tween for 2h at room temperature. Staining of the treated cells with ab692 (5 µg/ml) was performed overnight at 4°C in PBS containing 1% BSA and 0.1% tween. A **anti-mouse DyLight 488** polyclonal antibody (**ab96879**) at 1/250 dilution was used as the secondary antibody. Nuclei were counterstained with DAPI and are shown in blue.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Bcl-2 antibody [100/D5] (ab692)

Immunohistochemical analysis of paraffin-embedded human tonsil tissue labeling Bcl-2 with ab692 at 1/100 dilution. Samples were incubated with primary antibody for 30-45 minutes at RT.



Flow Cytometry - Anti-Bcl-2 antibody [100/D5] (ab692)

Overlay histogram showing Jurkat cells stained with ab692 (red line). The cells were fixed with 4% paraformaldehyde (10 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab692, 1/10 dilution) for 30 min at 22°C. The secondary antibody used was a goat **anti-mouse DyLight® 488** (IgG; H+L) (**ab96879**) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was mouse IgG1 [ICIGG1] (**ab91353**, 2µg/1x10<sup>6</sup> cells) used under the same conditions. Acquisition of >5,000 events was performed. This antibody gave a positive signal in Jurkat cells fixed with methanol (5 min)/permeabilized in 0.1% PBS-Tween used under the same conditions.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

#### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors