


Anti-beta Actin antibody [mAbcam 8224] - Loading Control ab8224

★★★★★ [35 Abreviews](#) [472 References](#) [画像数 10](#)

製品の概要

製品名	Anti-beta Actin antibody [mAbcam 8224] - Loading Control
製品の詳細	Mouse monoclonal [mAbcam 8224] to beta Actin - Loading Control
由来種	Mouse
特異性	Recognises a single band at 42kD representing beta Actin. The immunogen used for this product shares 77% homology with Gamma actin/actin cytoplasmic 2. Cross-reactivity with this protein has not been confirmed experimentally.
アプリケーション	適用あり: WB, IHC-P, ICC/IF 適用なし: Flow Cyt (Intra)
種交差性	交差種: Mouse, Rat, Human, Xenopus laevis, Drosophila melanogaster, Schizosaccharomyces pombe 交差が予測される動物種: Rabbit, Chicken, Cow, Cat, Dog, Pig, Chinese hamster, Other species 
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers. (Peptide available as ab13772)
ポジティブ・コントロール	WB: A431; HEK293; NIH3T3; PC12 whole cell lysates; Xenopus embryo lysate; Drosophila lysate; S. pombe lysate. ICC/IF: HeLa cells. IHC/P: Human colon (FFPE)
特記事項	<p>This monoclonal antibody to beta actin works well as a protein loading control in Western blot for a broad range of species including Xenopus, Drosophila and S. pombe.</p> <p>This antibody clone [mAbcam 8224] is manufactured by Abcam.</p> <p>If you require this antibody in a particular buffer formulation or a particular conjugate for your experiments, please contact orders@abcam.com or you can find further information here.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	pH: 7.40 Preservative: 0.02% Sodium azide Constituents: PBS, 6.97% L-Arginine
精製度	Protein G purified
一次抗体 備考	This clone works well as a loading control for Xenopus, Drosophila, S. cerevisiae and S.pombe. We recommend using ab8224 instead of ab8226 for these species.
ポリ/モノ	モノクローナル
クローン名	mAbcam 8224
ミエローマ	Sp2/0-Ag14
アイソタイプ	IgG1
軽鎖の種類	kappa

アプリケーション

The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab8224の使用に適用されます**

アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB	★★★★★ (22)	Use a concentration of 1 µg/ml. Detects a band of approximately 42 kDa (predicted molecular weight: 42 kDa). Can be blocked with Human beta Actin peptide (ab13772) . This antibody has been designed for use as a loading control and is ideal for this purpose. Block membrane for 1 hr in 5%BSA. Incubate antibody in TBST for one hour or more.
IHC-P	★★★★★ (7)	Use a concentration of 1 µg/ml. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
ICC/IF	★★★★★ (6)	Use a concentration of 1 µg/ml.

追加情報 Is unsuitable for Flow Cyt (Intra).

ターゲット情報

機能	Actins are highly conserved proteins that are involved in various types of cell motility and are ubiquitously expressed in all eukaryotic cells.
関連疾患	Defects in ACTB are a cause of dystonia juvenile-onset (DYTJ) [MIM:607371]. DYTJ is a form of dystonia with juvenile onset. Dystonia is defined by the presence of sustained involuntary muscle contraction, often leading to abnormal postures. DYTJ patients manifest progressive, generalized, dopa-unresponsive dystonia, developmental malformations and sensory hearing loss.
配列類似性	Belongs to the actin family.


翻訳後修飾

ISGylated.

細胞内局在

Cytoplasm > cytoskeleton. Localized in cytoplasmic mRNP granules containing untranslated mRNAs.

画像



All lanes : Anti-beta Actin antibody [mAbcam 8224] - Loading Control (ab8224) at 1 µg/ml

Lane 1 : A431 (Human epithelial carcinoma cell line) Whole Cell Lysate

Lane 2 : HEK293 (Human embryonic kidney cell line) Whole Cell Lysate

Lane 3 : NIH 3T3 (Mouse embryonic fibroblast cell line) Whole Cell Lysate

Lane 4 : PC12 (Rat adrenal pheochromocytoma cell line) Whole Cell Lysate

Lane 5 : Skeletal Muscle (Human) Tissue Lysate - adult normal tissue

Lysates/proteins at 20 µg per lane.


Secondary

All lanes : Goat Anti-Mouse IgG H&L (Alexa Fluor® 790) ([ab175783](#)) at 1/10000 dilution

Predicted band size: 42 kDa

Observed band size: 42 kDa

This blot was produced using a 4-12% Bis-tris gel under the MOPS buffer system. The gel was run at 200V for 50 minutes before being transferred onto a Nitrocellulose membrane at 30V for 70 minutes. The membrane was then blocked for an hour using 5% Milk before being incubated with ab8224 overnight at 4°C. Antibody binding was detected using a goat [anti-mouse Alexa Fluor 790](#) ([ab175783](#)) at a 1:10,000 dilution for 1hr at room temperature and then imaged using the Licor Odyssey CLx.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-beta Actin antibody [mAbcam 8224] - Loading Control (ab8224)

IHC image of ab8224 staining beta Actin in human colon formalin fixed paraffin embedded tissue sections*, performed on a Leica Bond. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab8224, 1µg/ml working concentration, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX. No primary antibody was used in the secondary only control (shown on the inset).


For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.

*Tissue obtained from the Human Research Tissue Bank, supported by the NIHR Cambridge Biomedical Research Centre



Immunocytochemistry/ Immunofluorescence - Anti-beta Actin antibody [mAbcam 8224] - Loading Control (ab8224)

ab8224 staining beta Actin in HeLa cells. The cells were fixed with 100% methanol (5 min), permeabilized with 0.1% PBS-Tween for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1%PBS-Tween for 1h. The cells were then incubated overnight at 4°C with ab8224 at 1µg/ml and **ab6046**, Rabbit polyclonal to beta Tubulin - Loading Control. Cells were then incubated with **ab150117**, Goat polyclonal Secondary Antibody to Mouse IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 dilution (shown in green) and **ab150080**, Goat polyclonal Secondary Antibody to Rabbit IgG - H&L (Alexa Fluor® 594) at 1/1000 dilution (shown in pseudocolour magenta). Nuclear DNA was labelled with DAPI (shown in blue). Image was acquired with a high-content analyser (Operetta CLS, Perkin Elmer) and a maximum intensity projection of confocal sections is shown.



Western blot - Anti-beta Actin antibody [mAbcam 8224] - Loading Control (ab8224)

All lanes : Anti-beta Actin antibody [mAbcam 8224] - Loading Control (ab8224) at 1 µg/ml

Lane 1 : HeLa (Human epithelial carcinoma cell line) Whole Cell Lysate

Lane 2 : NIH 3T3 (Mouse embryonic fibroblast cell line) Whole Cell Lysate

Lane 3 : PC12 (Rat adrenal pheochromocytoma cell line) Whole Cell Lysate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : Goat Anti-Mouse IgG H&L (HRP) preadsorbed (**ab97040**) at 1/50000 dilution


Performed under reducing conditions.

Predicted band size: 42 kDa

Observed band size: 42 kDa

Exposure time: 3 minutes

This blot was produced using a 4-12% Bis-tris gel under the MOPS buffer system. The gel was run at 200V for 50 minutes before being transferred onto a Nitrocellulose membrane at 30V for 70 minutes. The membrane was then blocked for an hour using 2% Bovine Serum Albumin before being incubated with ab8224 overnight at 4°C. Antibody binding was detected using an anti-mouse antibody conjugated to HRP, and visualised using ECL development solution **ab133406**




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-beta Actin antibody [mAbcam 8224] - Loading Control (ab8224)

IHC image of beta actin staining in human colon formalin fixed paraffin embedded tissue section*. The section was pre-treated using pressure cooker heat mediated antigen retrieval with sodium citrate buffer (pH6) for 30mins. The section was incubated with ab8224, 3µg/ml overnight at +4°C. A goat **anti-mouse HRP**-conjugated secondary antibody (**ab6789**, 1/2000 dilution) was used for 1hr at room temperature. The section was counterstained with haematoxylin and mounted with DPX.

The inset negative control image is secondary-only at 1/500 dilution.

*Tissue obtained from the Human Research Tissue Bank, supported by the NIHR Cambridge Biomedical Research Centre



Western blot - Anti-beta Actin antibody [mAbcam 8224] - Loading Control (ab8224)

Anti-beta Actin antibody [mAbcam 8224] - Loading Control (ab8224) + Xenopus embryo lysate at 20 µg

Secondary

Rabbit Anti-Mouse IgG H&L (HRP) (**ab6728**)

Developed using the ECL technique.

Performed under reducing conditions.

Predicted band size: 42 kDa


ab8224 used on Xenopus embryo lysate (20 ug of lysate/lane).

Secondary

Rabbit polyclonal **anti-mouse HRP** was used as the secondary antibody (**ab6728**) and developed using the ECL technique.

Performed under reducing conditions.

Predicted band size : 42kD



Western blot - Anti-beta Actin antibody [mAbcam 8224] - Loading Control (ab8224)

All lanes : Anti-beta Actin antibody [mAbcam 8224] - Loading Control (ab8224) at 1 µg/ml

Lane 1 : Drosophila lysate

Lane 2 : *S. pombe* lysate

Lane 3 : *S. cerevisiae* lysate (Actin 1 - please see note)

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Rabbit Anti-Mouse IgG H&L (HRP) ([ab6728](#)) at 1/5000 dilution

Developed using the ECL technique.


Performed under reducing conditions.

Predicted band size: 42 kDa

Observed band size: 42 kDa

Note: although *S. cerevisiae* is not known to express beta Actin, Abcam believes that the band on lane 3 corresponds to Actin 1 (Swissprot ID: P60010, based on sequence similarity).

Secondary antibody - rabbit **anti-mouse HRP** ([ab6728](#))



Western blot - Anti-beta Actin antibody [mAbcam 8224] - Loading Control (ab8224)

This image is courtesy of an anonymous Abreview

All lanes : Anti-beta Actin antibody [mAbcam 8224] - Loading Control (ab8224) at 1/1000 dilution

Lane 1 : Fruit fly (*Drosophila melanogaster*) whole cell lysate - Female

Lane 2 : Fruit fly (*Drosophila melanogaster*) whole cell lysate - Male

Lysates/proteins at 100 µg per lane.

Secondary

All lanes : An HRP-conjugated Sheep polyclonal to mouse IgG at 1/10000 dilution

Developed using the ECL technique.

Performed under reducing conditions.

Predicted band size: 42 kDa

Exposure time: 2 minutes

Blocking step: 5% Milk for 1 hour at 20°C.

Anti-beta Actin antibody [mAbcam 8224] - Loading Control (ab8224) at 1/10000 dilution + Mouse CT26 cells at 40 µg


Secondary

Anti-mouse IgG, HRP-linked Antibody at 1/10000 dilution

Developed using the ECL technique.

Predicted band size: 42 kDa

Observed band size: 42 kDa




Western blot - Anti-beta Actin antibody [mAbcam 8224] - Loading Control (ab8224)

This image is courtesy of an Abreview submitted by Joel Ohana

Exposure time: 6 minutes

Western blot analysis using ab8224 at 1:1000 on Mouse CT26 cells. Blocking agent and dilution buffer was 5% milk in TBST



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-beta Actin antibody

[mAbcam 8224] - Loading Control (ab8224)

This image is courtesy of an abreview submitted by Carl Hobbs, King's College London, United Kingdom

Immunohistochemical detection of beta Actin using antibody [mAbcam 8224] - Loading Control on formaldehyde-fixed paraffin-embedded rat cerebellum sections. Antigen retrieval step: heat mediated Citric acid pH6 buffer. Permeabilization: No. Blocking step: 1% BSA for 10 mins @ rt°C. Primary antibody dilution 1/1000 for 2 hours in TBS/BSA/azide. Secondary Antibody: anti Mouse Igs conjugated to biotin (1/200). beta Actin appears to be particularly enriched not only in the glomeruli of the Granule cell layer (indicated by red arrowheads) but also in Microglia (indicated by green arrowheads); All positive microglia appear to be ramified thus not presumed to be activated.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors