


# Biotin Anti-CD3 antibody [UCHT1] ab191112

[1 References](#) [画像数 2](#)

### 製品の概要

製品名	Biotin Anti-CD3 antibody [UCHT1]
製品の詳細	Biotin Mouse monoclonal [UCHT1] to CD3
由来種	Mouse
標識	Biotin
特異性	ab191112 recognizes the CD3 antigen of the TCR/CD3 complex on mature Human T cells. The antibody reacts with the epsilon chain of the CD3 complex.
アプリケーション	<b>適用あり:</b> Flow Cyt (Intra), Flow Cyt
種交差性	<b>交差種:</b> Human <b>交差が予測される動物種:</b> Non human primates 
免疫原	Tissue, cells or virus corresponding to Human CD3. Human thymocytes followed by Sezary T cells.
ポジティブ・コントロール	Flow Cytometry: human peripheral whole blood. Flow Cyt (Intra): Human blood.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle. Store In the Dark.
バッファー	pH: 7.40 Preservative: 0.0975% Sodium azide Constituent: 99% PBS
精製度	Size exclusion
特記事項 (精製)	ab191112 is free of unconjugated biotin.


ポリ/モノ	モノクローナル
クローン名	UCHT1
アイソタイプ	IgG1

## アプリケーション


**The Abpromise guarantee** **Abpromise保証は、**次のテスト済みアプリケーションにおけるab191112の使用に適用されます  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
Flow Cyt (Intra)		Use a concentration of 1 - 12 µg/ml.
Flow Cyt		Use a concentration of 1 - 12 µg/ml.

## 画像



Flow cytometry surface staining pattern of human peripheral whole blood stained using ab191112 (concentration in sample 0.3 µg/mL) streptavidin APC.



Separation of human CD3 positive lymphocytes (red-filled) from neutrophil granulocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using ab191112 (concentration in sample 0.3 µg/mL) streptavidin APC.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors