

### Biotin Anti-Plasminogen antibody ab7335

★★★★★ [1 Abreviews](#) [1 References](#)

#### 製品の概要

製品名	Biotin Anti-Plasminogen antibody
製品の詳細	Biotin Goat polyclonal to Plasminogen
由来種	Goat
標識	Biotin
アプリケーション	<b>適用あり:</b> Flow Cyt, ELISA, WB, ICC/IF
種交差性	<b>交差種:</b> Human
免疫原	Full length native protein (purified) corresponding to Human Plasminogen. Human plasma Database link: <a href="#">P00747</a>
特記事項	<p>Biotinamidocaproate N-Hydroxysuccinimide Ester (BAC) Biotin/Protein Ratio: 10-20 BAC molecules per goat IgG molecule.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

#### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	<p>pH: 6.50</p> <p>Preservative: 0.01% Sodium azide</p> <p>Constituents: 0.88% Sodium chloride, 1% BSA, 0.42% Tripotassium orthophosphate</p>
精製度	IgG fraction
特記事項 (精製)	IgG fraction antibody purified from monospecific antiserum by a multi-step process including delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer.
ポリ/モノ	ポリクローナル

## アプリケーション

**The Abpromise guarantee** Abpromise保証は、次のテスト済みアプリケーションにおけるab7335の使用に適用されます  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
Flow Cyt	★★★★★ (1)	Use at an assay dependent concentration. <b>ab37376</b> - Goat polyclonal IgG, is suitable for use as an isotype control with this antibody.
ELISA		Use at an assay dependent concentration. As well as other antibody based assays using streptavidin or avidin conjugates requiring lot-to-lot consistency.
WB		Use at an assay dependent concentration.
ICC/IF		Use at an assay dependent concentration. PubMed: 19182206

## ターゲット情報

## 機能

Plasmin dissolves the fibrin of blood clots and acts as a proteolytic factor in a variety of other processes including embryonic development, tissue remodeling, tumor invasion, and inflammation. In ovulation, weakens the walls of the Graafian follicle. It activates the urokinase-type plasminogen activator, collagenases and several complement zymogens, such as C1 and C5. Cleavage of fibronectin and laminin leads to cell detachment and apoptosis. Also cleaves fibrin, thrombospondin and von Willebrand factor. Its role in tissue remodeling and tumor invasion may be modulated by CSPG4. Binds to cells.

Angiostatin is an angiogenesis inhibitor that blocks neovascularization and growth of experimental primary and metastatic tumors in vivo.

## 組織特異性

Present in plasma and many other extracellular fluids. It is synthesized in the liver.

## 関連疾患

Defects in PLG are a cause of susceptibility to thrombosis (THR) [MIM:188050]. It is a multifactorial disorder of hemostasis characterized by abnormal platelet aggregation in response to various agents and recurrent thrombi formation.

Defects in PLG are the cause of plasminogen deficiency (PLGD) [MIM:217090]. PLGD is characterized by decreased serum plasminogen activity. Two forms of the disorder are distinguished: type 1 deficiency is additionally characterized by decreased plasminogen antigen levels and clinical symptoms, whereas type 2 deficiency, also known as dysplasminogenemia, is characterized by normal, or slightly reduced antigen levels, and absence of clinical manifestations. Plasminogen deficiency type 1 results in markedly impaired extracellular fibrinolysis and chronic mucosal pseudomembranous lesions due to subepithelial fibrin deposition and inflammation. The most common clinical manifestation of type 1 deficiency is ligneous conjunctivitis in which pseudomembranes formation on the palpebral surfaces of the eye progresses to white, yellow-white, or red thick masses with a wood-like consistency that replace the normal mucosa.

## 配列類似性

Belongs to the peptidase S1 family. Plasminogen subfamily.  
Contains 5 kringle domains.

	Contains 1 PAN domain. Contains 1 peptidase S1 domain.
<b>ドメイン</b>	Kringle domains mediate interaction with CSPG4.
<b>翻訳後修飾</b>	N-linked glycan contains N-acetylglucosamine and sialic acid. O-linked glycans consist of Gal-GalNAc disaccharide modified with up to 2 sialic acid residues (microheterogeneity). In the presence of the inhibitor, the activation involves only cleavage after Arg-580, yielding two chains held together by two disulfide bonds. In the absence of the inhibitor, the activation involves additionally the removal of the activation peptide.
<b>細胞内局在</b>	Secreted. Locates to the cell surface where it is proteolytically cleaved to produce the active plasmin. Interaction with HRG tethers it to the cell surface.
<b>製品の状態</b>	Cleaved into the following 5 chains: 1. Plasmin heavy chain A2. Activation peptide3. Angiostatin4. Plasmin heavy chain A, short form5. Plasmin light chain B

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors