

Biotin Rat IgM, kappa monoclonal [RTK2118] - Isotype control ab35773

製品の概要

製品名	Biotin Rat IgM, kappa monoclonal [RTK2118] - Isotype control
標識	Biotin
アプリケーション	適用あり: IHC-Fr, IHC-P, WB, IP, Flow Cyt
特記事項	<p>Isotype Control monoclonal antibodies are used to estimate the non-specific binding of target primary antibodies to cell surface antigens. Non-specific binding is due to Fc receptor binding or other protein-protein interactions. Isotype controls show negligible cross-reactivity with cell surface antigens on tissue sections or in cellular preparations. Isotype Controls should be used at identical concentrations and staining conditions as the target primary antibodies.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C. Do Not Freeze.
バッファー	<p>pH: 7.20</p> <p>Preservative: 0.09% Sodium azide</p> <p>Constituent: PBS</p>
精製度	Purified IgM
アイソタイプ・コントロール 備考	Isotype Control monoclonal antibodies are used to estimate the non-specific binding of target primary antibodies to cell surface antigens. Non-specific binding is due to Fc receptor binding or other protein-protein interactions. Isotype controls show negligible cross-reactivity with cell surface antigens on tissue sections or in cellular preparations. Isotype Controls should be used at identical concentrations and staining conditions as the target primary antibodies.

ポリ/モノ	モノクローナル
クローン名	RTK2118
アイソタイプ	IgM
軽鎖の種類	kappa

アプリケーション

The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab35773の使用に適用されます**
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-Fr		Use at an assay dependent dilution.
IHC-P		Use at an assay dependent dilution.
WB		Use at an assay dependent dilution.
IP		Use at an assay dependent dilution.
Flow Cyt		Use at an assay dependent dilution.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors