

Biotin Anti-Toxoplasma gondii antibody ab69251

製品の概要

製品名	Biotin Anti-Toxoplasma gondii antibody
製品の詳細	Biotin Goat polyclonal to Toxoplasma gondii
由来種	Goat
標識	Biotin
アプリケーション	適用あり: ELISA
種交差性	交差種: Toxoplasma gondii
免疫原	Tissue, cells or virus corresponding to Toxoplasma gondii Toxoplasma gondii. RH Strain of Toxoplasma gondii.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	<p>pH: 7.2</p> <p>Preservative: 0.1% Sodium azide</p> <p>Constituent: 0.0268% PBS</p>
精製度	Protein G purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee

Abpromise保証は、次のテスト済みアプリケーションにおけるab69251の使用に適用されます

アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
ELISA		1/1000.

ターゲット情報

関連性	<p>Toxoplasma is a crescent shaped sporozoan that lives as an intracellular parasite in various tissues of many vertebrates and completes its life cycle in a single host. Its life cycle includes two phases called the intestinal (or enteroepithelial) and extraintestinal phases. The intestinal phase produces oocysts and occurs only in cats, wild as well as domesticated. The extraintestinal phase occurs in all infected animals including cats, and produces tachyzoites (actively proliferating trophozoites) and eventually, bradyzoites (slowly growing trophozoites) or zoitocysts. Infection due to Toxoplasma gondii occurs in pregnant women where a variable degree of immunosuppression may exist or in patients receiving immunosuppressive drug therapy. Toxoplasma infects tissue of the GI tract where an active infection is accompanied by fever and enlargement of the spleen. Symptoms of toxoplasmosis are generally mild but severe infection of lymph nodes may occur. Congenital toxoplasmosis, in which the maternal infection is transmitted during pregnancy, can produce blindness or mental retardation in the newborn.</p>
-----	---

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors