

### Biotin Anti-TRAIL antibody [N2B2] ab95531

[2 References](#) [画像数 1](#)

#### 製品の概要

---

製品名	Biotin Anti-TRAIL antibody [N2B2]
製品の詳細	Biotin Rat monoclonal [N2B2] to TRAIL
由来種	Rat
標識	Biotin
アプリケーション	<b>適用あり:</b> Flow Cyt
種交差性	<b>交差種:</b> Recombinant fragment
免疫原	Tissue, cells or virus corresponding to Mouse TRAIL. Mouse TRAIL-transfected 2PK-3 cells.
ポジティブ・コントロール	Mouse TRAIL transfected cells
特記事項	<p>ab95531 inhibits IL2- or IL15-activated NK cell cytotoxicity against Mouse fibrosarcoma L929 target cells.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

#### 製品の特性

---

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	pH: 7.20 Preservative: 0.09% Sodium azide Constituents: 0.87% Sodium chloride, PBS
精製度	Protein G purified
一次抗体 備考	ab95531 inhibits IL2- or IL15-activated NK cell cytotoxicity against Mouse fibrosarcoma L929 target cells.

ポリ/モノ	モノクローナル
クローン名	N2B2
アイソタイプ	IgG2a
軽鎖の種類	kappa

## アプリケーション


**The Abpromise guarantee** Abpromise保証は、次のテスト済みアプリケーションにおけるab95531の使用に適用されます  
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
Flow Cyt		Use 0.125-0.25µg for 10 <sup>5-8</sup> cells. in a final volume of 100 µL.  ab19445 Rat monoclonal IgG2a is suitable for use as an

## ターゲット情報

<b>機能</b>	Cytokine that binds to TNFRSF10A/TRAILR1, TNFRSF10B/TRAILR2, TNFRSF10C/TRAILR3, TNFRSF10D/TRAILR4 and possibly also to TNFRSF11B/OPG. Induces apoptosis. Its activity may be modulated by binding to the decoy receptors TNFRSF10C/TRAILR3, TNFRSF10D/TRAILR4 and TNFRSF11B/OPG that cannot induce apoptosis.
<b>組織特異性</b>	Widespread; most predominant in spleen, lung and prostate.
<b>配列類似性</b>	Belongs to the tumor necrosis factor family.
<b>細胞内局在</b>	Membrane.

## 画像



Flow Cytometry - Biotin Anti-TRAIL antibody [N2B2]  
(ab95531)

ab95531, at 0.125 µg, staining of TRAIL transfected cells (filled) or with Rat IgG2a Isotype Control (open) followed by Streptavidin PE by Flow Cytometry. Total viable cells were used for analysis.

## **Our Abpromise to you: Quality guaranteed and expert technical support**

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

## **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors