

Anti-BNIP3 antibody [EPR4034] ab109362

KO 評価済 リコンビナント RabMAb

★★★★☆ 3 Abreviews 62 References 画像数 9

製品の概要

| | |
|--------------|---|
| 製品名 | Anti-BNIP3 antibody [EPR4034] |
| 製品の詳細 | Rabbit monoclonal [EPR4034] to BNIP3 |
| 由来種 | Rabbit |
| アプリケーション | 適用あり: WB, IHC-P, ICC/IF |
| 種交差性 | 交差種: Mouse, Rat, Human |
| 免疫原 | Synthetic peptide. This information is proprietary to Abcam and/or its suppliers. |
| ポジティブ・コントロール | WB: Jurkat cells, Jurkat cells treated with etoposide, SH-SY5Y cells, SH-SY5Y cells treated with camptothecin and MCF-7 cells treated with Cocl2 lysates and rat kidney, mouse spleen and mouse kidney tissue lysates. IHC-P: Human renal adenocarcinoma and kidney tissues. ICC/IF: HeLa and SH-SY5Y cells. |
| 特記事項 | <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p> |

製品の特性

| | |
|-------|---|
| 製品の状態 | Liquid |
| 保存方法 | Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Stable for 12 months at -20°C. |
| バッファー | <p>pH: 7.20</p> <p>Preservative: 0.01% Sodium azide</p> <p>Constituents: 40% Glycerol, 59% PBS, 0.05% BSA</p> |
| 精製度 | Protein A purified |
| ポリ/モノ | モノクローナル |
| クローン名 | EPR4034 |

アプリケーション


The Abpromise guarantee Abpromise保証は、 次のテスト済みアプリケーションにおけるab109362の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

| アプリケーション | Abreviews | 特記事項 |
|----------|-----------|---|
| WB | ★★★★★ (3) | 1/1000. Detects a band of approximately 30 kDa (predicted molecular weight: 22 kDa). For unpurified use at 1/1000 - 1/10000. |
| IHC-P | | 1/100 - 1/250. Perform heat mediated antigen retrieval before commencing with IHC staining protocol. See <u>IHC antigen retrieval protocols.</u> (Heat to 98°C, allow to cool for 10-20 minutes). For unpurified use at 1/50 - 1/100. <u>ab172730</u> - Rabbit monoclonal IgG, is suitable for use as an |
| ICC/IF | | 1/150. For unpurified use at 1/100 - 1/250. |

ターゲット情報

| | |
|-------|--|
| 機能 | Apoptosis-inducing protein that, which can overcome BCL2 suppression. May play a role in repartitioning calcium between the two major intracellular calcium stores in association with BCL2. |
| 配列類似性 | Belongs to the NIP3 family. |
| 細胞内局在 | Mitochondrion. Mitochondrion membrane. Coexpression with the E1B 19-kDa protein results in a shift in NIP3 localization pattern to the nuclear envelope. Colocalizes with ACAA2 in the mitochondria. |

画像



Lane 1: Wild-type HAP1 whole cell lysate (40 µg)


Lane 2: BNIP3 knockout HAP1 whole cell lysate (40 µg)

Lane 3: SHSY5Y whole cell lysate (40 µg)

Lane 4: A431 whole cell lysate (40 µg)

Lanes 1 - 4: Merged signal (red and green). Green - ab109362 observed at 25 kDa. Red - loading control, **ab9484**, observed at 37 kDa.

ab109362 was shown to recognize BNIP3 in wild-type HAP1 cells as signal was lost at the expected MW in BNIP3 knockout cells. Additional cross-reactive bands were observed in the wild-type and knockout cells. Wild-type and BNIP3 knockout samples were subjected to SDS-PAGE. ab109362 and **ab9484** (Mouse anti-GAPDH loading control) were incubated overnight at 4°C at 1/1000 dilution and 1/20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed **ab216773** and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed **ab216776** secondary antibodies at 1/20000 dilution for 1 hour at room temperature before imaging.



Anti-BNIP3 antibody [EPR4034] (ab109362) at 1/1000 dilution (purified) + SH-SY5Y cell lysate at 20 µg


Secondary

Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

Predicted band size: 22 kDa

Observed band size: 30 kDa

Blocking and diluting buffer: 5% NFDM/TBST.



Western blot - Anti-BNIP3 antibody [EPR4034]
(ab109362)

Anti-BNIP3 antibody [EPR4034] (ab109362) at 1/2000 dilution
(purified) + Jurkat cell lysate at 10 µg


Secondary

Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000
dilution

Predicted band size: 22 kDa

Observed band size: 30 kDa

Blocking and diluting buffer: 5% NFDM/TBST.



Western blot - Anti-BNIP3 antibody [EPR4034]
(ab109362)

All lanes : Anti-BNIP3 antibody [EPR4034] (ab109362) at 1/2000
dilution (purified)

Lane 1 : Rat kidney tissue lysate

Lane 2 : Mouse kidney tissue lysate

Lane 3 : Mouse spleen tissue lysate

Lysates/proteins at 20 µg per lane.


Secondary

All lanes : Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at
1/1000 dilution

Predicted band size: 22 kDa

Observed band size: 30 kDa

Blocking and diluting buffer: 5% NFDM/TBST.



Western blot - Anti-BNIP3 antibody [EPR4034]
(ab109362)

All lanes : Anti-BNIP3 antibody [EPR4034] (ab109362) at 1/1000 dilution (unpurified)

Lane 1 : Jurkat cell lysate treated with etoposide

Lane 2 : Jurkat cell lysate

Lane 3 : SH-SY5Y cell lysate treated with camptothecin


Lane 4 : SH-SY5Y cell lysate

Lane 5 : MCF-7 cell lysate treated with Cocl2

Lysates/proteins at 10 µg per lane.


Predicted band size: 22 kDa

Observed band size: 30 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-BNIP3 antibody
[EPR4034] (ab109362)


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human renal adenocarcinoma tissue labelling BNIP3 with purified ab109362 at 1/100. Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9. [ab97051](#), a HRP-conjugated goat anti-rabbit IgG (H+L) was used as the secondary antibody (1/500). Negative control using PBS instead of primary antibody. Counterstained with hematoxylin.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-BNIP3 antibody [EPR4034] (ab109362)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human kidney tissue labelling BNIP3 with unpurified ab109362 at a 1/50 dilution.

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunocytochemistry/ Immunofluorescence - Anti-BNIP3 antibody [EPR4034] (ab109362)

Immunocytochemistry/Immunofluorescence analysis of SH-SY5Y cells labelling BNIP3 with unpurified ab109362 at a 1/100 dilution.

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-BNIP3 antibody [EPR4034] (ab109362)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors