# abcam

### Product datasheet

## Anti-BRD8/p120 antibody ab17969

2 References 画像数 3

製品の概要

製品名 Anti-BRD8/p120 antibody

製品の詳細 Rabbit polyclonal to BRD8/p120

由来種 Rabbit

特異性 Specific to p120

アプリケーション 適用あり: IP, IHC-P, WB

種交差性 交差種: Human

交差が予測される動物種: Rat, Rabbit, Horse, Guinea pig, Dog, Pig, Rhesus monkey, Gorilla,

Opossum, Chinese hamster, Elephant

免疫原 Synthetic peptide within BRD8/p120 aa 200-300. The exact immunogen sequence used to

generate this antibody is proprietary information. If additional detail on the immunogen is needed to determine the suitability of the antibody for your needs, please **contact** our Scientific Support

team to discuss your requirements.

Database link: **Q9H0E9** 

ポジティブ・コントロール Nuclear extract from HeLa cells.

特記事項 The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or

contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

**バッファー** pH: 7

Preservative: 0.1% Sodium azide

Constituents: 0.021% PBS, 1.764% Sodium citrate, 1.815% Tris

精製度 Immunogen affinity purified

特記事項(精製) Antibody was affinity purified using an epitope specific to BRD8/p120 immobilized on solid

1

support.

**ポリ/モノ** ポリクローナル

アイソタイプ lgG

#### アプリケーション

The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab17969の使用に適用されますアプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IP		Use at an assay dependent concentration.
IHC-P		1/500 - 1/2000. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
WB		1/10000 - 1/25000. Detects a band of approximately 170 kDa (predicted molecular weight: 138 kDa).

#### ターゲット情報

#### 機能

May act as a coactivator during transcriptional activation by hormone-activated nuclear receptors (NR). Isoform 2 stimulates transcriptional activation by AR/DHTR, ESR1/NR3A1, RXRA/NR2B1 and THRB/ERBA2. At least isoform 1 and isoform 2 are components of the NuA4 histone acetyltransferase (HAT) complex which is involved in transcriptional activation of select genes principally by acetylation of nucleosomal histones H4 and H2A. This modification may both alter nucleosome - DNA interactions and promote interaction of the modified histones with other proteins which positively regulate transcription. This complex may be required for the activation of transcriptional programs associated with oncogene and proto-oncogene mediated growth induction, tumor suppressor mediated growth arrest and replicative senescence, apoptosis, and DNA repair. NuA4 may also play a direct role in DNA repair when recruited to sites of DNA damage. Component of a SWR1-like complex that specifically mediates the removal of histone H2A.Z/H2AFZ from the nucleosome.

#### 組織特異性

Expressed in adipose tissue, brain, heart, kidney, liver, lung, pancreas, placenta and skeletal

muscle.


配列類似性

Contains 2 bromo domains.

細胞内局在

Nucleus.

#### 画像




Immunoprecipitation - Anti-BRD8/p120 antibody (ab17969)

ab17969 immunoprecipitating BRD8 at 6 µg per reaction.

**Lane 1:** ab17969 in HeLa nuclear extract (0.5 or 1 mg per IP reaction; 20% of IP loaded).


Lane 2: IgG control

For blotting immunoprecipitated BRD8, ab17969 was used at 1  $\mu$ g/mL. Detection: Chemiluminescence with an exposure time of 3 minutes.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-BRD8/p120 antibody (ab17969)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human prostate carcinoma tissue labelling BRD8 / p120 with ab17969 at 1/1000 (1µg/ml). Detection: DAB.



Detection of human BRD8 by Western Blot. Sample is nuclear extract (NE; 5, 15, and 50mcg) from HeLa cells.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

#### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- · Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="https://www.abcam.co.jp/abpromise">https://www.abcam.co.jp/abpromise</a> or contact our technical team.

### Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors