


### Anti-Caveolin-1 antibody - Caveolae Marker ab18199

KO 評価済

★★★★★ [9 Abreviews](#) [27 References](#) [画像数 5](#)

#### 製品の概要

製品名	Anti-Caveolin-1 antibody - Caveolae Marker
製品の詳細	Rabbit polyclonal to Caveolin-1 - Caveolae Marker
由来種	Rabbit
アプリケーション	適用あり: IP, WB 適用なし: ICC/IF
種交差性	交差種: Mouse, Rat, Human, Recombinant fragment 交差が予測される動物種: Cow, Dog, Pig 
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

#### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	pH: 7.40 Preservative: 0.02% Sodium azide Constituents: PBS, 1% BSA  Batches of this product that have a concentration < 1mg/ml may have BSA added as a stabilising agent. If you would like information about the formulation of a specific lot, please contact our scientific support team who will be happy to help.
精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル

## アプリケーション

**The Abpromise guarantee** Abpromise保証は、次のテスト済みアプリケーションにおけるab18199の使用に適用されます  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IP		Use at an assay dependent concentration.
WB	★★★★★ (5)	Use a concentration of 1 µg/ml. Detects a band of approximately 20 kDa (predicted molecular weight: 20 kDa).


## 追加情報

Is unsuitable for ICC/IF.

## ターゲット情報

機能	May act as a scaffolding protein within caveolar membranes. Interacts directly with G-protein alpha subunits and can functionally regulate their activity (By similarity). Involved in the costimulatory signal essential for T-cell receptor (TCR)-mediated T-cell activation. Its binding to DPP4 induces T-cell proliferation and NF-kappa-B activation in a T-cell receptor/CD3-dependent manner. Recruits CTNNB1 to caveolar membranes and may regulate CTNNB1-mediated signaling through the Wnt pathway.
組織特異性	Expressed in muscle and lung, less so in liver, brain and kidney.
関連疾患	Defects in CAV1 are the cause of congenital generalized lipodystrophy type 3 (CGL3) [MIM:612526]; also called Berardinelli-Seip congenital lipodystrophy type 3 (BSCL3). Congenital generalized lipodystrophies are autosomal recessive disorders characterized by a near absence of adipose tissue, extreme insulin resistance, hypertriglyceridemia, hepatic steatosis and early onset of diabetes.
配列類似性	Belongs to the caveolin family.
翻訳後修飾	The initiator methionine for isoform Beta is removed during or just after translation. The new N-terminal amino acid is then N-acetylated.
細胞内局在	Golgi apparatus membrane. Cell membrane. Membrane > caveola. Membrane raft. Colocalized with DPP4 in membrane rafts. Potential hairpin-like structure in the membrane. Membrane protein of caveolae.

## 画像



**All lanes :** Anti-Caveolin-1 antibody - Caveolae Marker (ab18199) at 1 µg/ml

**Lane 1 :** Wild-type A-431 (Human epidermoid carcinoma cell line) whole cell lysate

**Lane 2 :** CAV1 knockout A-431 (Human epidermoid carcinoma cell line) whole cell lysate

**Lane 3 :** A549 (Human lung carcinoma cell line) whole cell lysate

**Lane 4 :** HeLa (Human epithelial cell line from cervix adenocarcinoma) whole cell lysate

Lysates/proteins at 20 µg per lane.


Performed under reducing conditions.

**Predicted band size:** 20 kDa

**Observed band size:** 21-24 kDa

**Lanes 1 -4:** Merged signal (red and green). Green - ab18199 observed at 21-24 kDa. Red - loading control, **ab7291** (Mouse anti-Alpha Tubulin [DM1A]) observed at 55kDa.

ab18199 was shown to react with Caveolin-1 in wild-type A431 cells in western blot. Loss of signal was observed when CAV1 knockout sample was used. Wild-type and CAV1 knockout A431 cell lysates were subjected to SDS-PAGE. Membranes were blocked in 3% milk in TBS-T (0.1% Tween®) before incubation with ab18199 and **ab7291** (Mouse anti-Alpha Tubulin [DM1A]) overnight at 4°C at 1 µg/ml and a 1 in 20000 dilution respectively. Blots were incubated with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed (**ab216776**) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.



Western blot - Anti-Caveolin-1 antibody - Caveolae Marker (ab18199)

**All lanes :** Anti-Caveolin-1 antibody - Caveolae Marker (ab18199) at 1 µg/ml

**Lane 1 :** Mouse Heart Lysate

**Lane 2 :** Human Heart Lysate

Lysates/proteins at 10 µg per lane.


#### Secondary

**All lanes :** Alexa Fluor Goat polyclonal to Rabbit IgG (700) at 1/5000 dilution

Performed under reducing conditions.

**Predicted band size:** 20 kDa

**Observed band size:** 20 kDa



Western blot - Anti-Caveolin-1 antibody - Caveolae Marker (ab18199)

**All lanes :** Anti-Caveolin-1 antibody - Caveolae Marker (ab18199) at 1 µg/ml

**Lane 1 :** NIH/3T3 whole cell lysate ([ab7179](#))

**Lane 2 :** MEF1 (Mouse embryonic fibroblast cell line) Whole Cell Lysate

**Lane 3 :** Heart (Mouse) Tissue Lysate

**Lane 4 :** Heart (Rat) Tissue Lysate

Lysates/proteins at 10 µg per lane.


#### Secondary

**All lanes :** IRDye 680 Conjugated Goat Anti-Rabbit IgG (H+L) at 1/10000 dilution

Performed under reducing conditions.

**Predicted band size:** 20 kDa

**Observed band size:** 20 kDa




Caveolin 1 - Caveolae Marker was immunoprecipitated using 0.5mg Mouse Heart whole tissue lysate, 5µg of Rabbit polyclonal to Caveolin 1 and 50µl of protein G magnetic beads (+). No antibody was added to the control (-).

The antibody was incubated under agitation with Protein G beads for 10min, Mouse Heart whole tissue lysate lysate diluted in RIPA buffer was added to each sample and incubated for a further 10min under agitation.

Proteins were eluted by addition of 40µl SDS loading buffer and incubated for 10min at 70°C; 10µl of each sample was separated on a SDS PAGE gel, transferred to a nitrocellulose membrane, blocked with 5% BSA and probed with ab18199.

Secondary: Clean blot (HRP conjugate) at 1/1000 dilution.

Band: 20kDa: Caveolin 1 - Caveolae Marker.



Anti-Caveolin-1 antibody - Caveolae Marker (ab18199) at 1 µg/ml + Recombinant Human Caveolin-1 protein ([ab114170](#)) at 0.01 µg

### Secondary

Goat Anti-Rabbit IgG H&L (HRP) preadsorbed ([ab97080](#)) at 1/5000 dilution

Developed using the ECL technique.

Performed under reducing conditions.

**Predicted band size:** 20 kDa

**Exposure time:** 1 minute

ab18199 recognizes the full length tagged recombinant Caveolin 1 protein ([ab114170](#)) which has an expected molecular weight of 46 kDa.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

**Our Abpromise to you: Quality guaranteed and expert technical support**

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

#### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors