abcam

Product datasheet

Anti-Caveolin-2 (phospho Y27) antibody [EP2221Y] ab76405

RabMAb

画像数1

製品の概要

製品名 Anti-Caveolin-2 (phospho Y27) antibody [EP2221Y]

製品の詳細 Rabbit monoclonal [EP2221Y] to Caveolin-2 (phospho Y27)

由来種 Rabbit

アプリケーション 適用あり: WB

適用なし: Flow Cyt,ICC/IF or IHC-P

種交差性 交差種: Human

免疫原 Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

ポジティブ・コントロール WB: HeLa cell lysates treated with pervanadate

特記事項 Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit

monoclonal antibodies. For details on our patents, please refer to $\underline{\textbf{RabMAb}}^{\underline{@}}\underline{\textbf{patents}}.$

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with these species. Please contact us for more information.

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

バッファー pH: 7.20

Preservative: 0.05% Sodium azide

Constituents: 0.1% BSA, 40% Glycerol (glycerin, glycerine), 9.85% Tris glycine, 50% Tissue

culture supernatant

精製度 Tissue culture supernatant

1

ポリ/モノ モノクローナル **ウローン名** EP2221Y

アイソタイプ IgG

アプリケーション

The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab76405の使用に適用されますアプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		1/1000 - 1/2000. Predicted molecular weight: 18 kDa.

追加情報 Is unsuitable for Flow Cyt,ICC/IF or IHC-P.

ターゲット情報

機能 May act as a scaffolding protein within caveolar membranes. Interacts directly with G-protein

alpha subunits and can functionally regulate their activity. Acts as an accessory protein in conjunction with CAV1 in targeting to lipid rafts and driving caveolae formation. The Ser-36 phosphorylated form has a role in modulating mitosis in endothelial cells. Positive regulator of cellular mitogenesis of the MAPK signaling pathway. Required for the insulin-stimulated nuclear translocation and activation of MAPK1 and STAT3, and the subsequent regulation of cell cycle

progression.

組織特異性 Expressed in endothelial cells, smooth muscle cells, skeletal myoblasts and fibroblasts.

配列類似性 Belongs to the caveolin family.

翻訳後修飾 Phosphorylated on serine and tyrosine residues. CAV1 promotes phosphorylation on Ser-23

which then targets the complex to the plasma membrane, lipid rafts and caveolae.

Phosphorylation on Ser-36 appears to modulate mitosis in endothelial cells (By similarity).

Phosphorylation on both Tyr-19 and Tyr-27 is required for insulin-induced 'Ser-727'

phosphorylation of STAT3 and its activation. Phosphorylation on Tyr-19 is required for insulininduced phosphorylation of MAPK1 and DNA binding of STAT3. Tyrosine phosphorylation is

induced by both EGF and insulin.


細胞内局在 Nucleus. Cytoplasm. Golgi apparatus membrane. Cell membrane. Membrane > caveola.

Potential hairpin-like structure in the membrane. Membrane protein of caveolae. Tyr-19-phosphorylated form is enriched at sites of cell-cell contact and is translocated to the nucleus in complex with MAPK1 in response to insulin (By similarity). Tyr-27-phosphorylated form is located

both in the cytoplasm and plasma membrane. CAV1-mediated Ser-23-phosphorylated form locates to the plasma membrane. Ser-36-phosphorylated form resides in intracellular

compartments.

画像



All lanes : Anti-Caveolin-2 (phospho Y27) antibody [EP2221Y] (ab76405) at 1/2000 dilution

Lane 1: HeLa cell lysates untreated.

Lane 2: HeLa cell lysates treated with pervanadate.

Lysates/proteins at 10 µg per lane.

Secondary

All lanes: HRP labelled goat anti-rabbit at 1/2000 dilution

Predicted band size: 18 kDa
Observed band size: 20 kDa

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors