abcam

Product datasheet

Anti-CD10 antibody ab73409

★★★★★ 2 Abreviews 4 References 画像数 3

製品の概要

製品名 Anti-CD10 antibody

製品の詳細 Rabbit polyclonal to CD10

由来種 Rabbit

特異性 From Jan 2024, QC testing of replenishment batches of this polyclonal changed. All tested and

expected application and reactive species combinations are still covered by our Abcam product promise. However, we no longer test all applications. For more information on a specific batch, please contact our Scientific Support who will be happy to help. You may also be interested in our

alternative recombinant antibody, ab256494.

アプリケーション 適用あり: WB, IHC-P, ICC/IF

種交差性 交差種: Human

交差が予測される動物種: Mouse, Rat, Rabbit, Cow 🔷

免疫原 Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

ポジティブ・コントロール This antibody gave a positive signal in the following lysates: LNCaP Nuclear; Ramos Whole Cell;

Raji Whole Cell; Human Kidney Tissue; Human Placenta Tissue This antibody gave a positive

result in IHC in the following FFPE tissue: human normal kidney.

特記事項
The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -

80°C. Avoid freeze / thaw cycle.

バッファー pH: 7.40

Preservative: 0.02% Sodium azide

Constituent: PBS

1

Batches of this product that have a concentration < 1mg/ml may have BSA added as a stabilising agent. If you would like information about the formulation of a specific lot, please contact our scientific support team who will be happy to help.

精製度 Immunogen affinity purified

ポリ/モノ ポリクローナル

アイソタイプ IgG

アプリケーション

The Abpromise guarantee <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab73409の使用に適用されますアプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		Use a concentration of 1 µg/ml. Detects a band of approximately 100 kDa (predicted molecular weight: 85 kDa).
IHC-P	★★ ☆☆☆(1)	Use a concentration of 1 µg/ml. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.
ICC/IF		Use a concentration of 5 µg/ml.

ターゲット情報

機能 Thermolysin-like specificity, but is almost confined on acting on polypeptides of up to 30 amino


acids. Biologically important in the destruction of opioid peptides such as Met- and Leuenkephalins by cleavage of a Gly-Phe bond. Able to cleave angiotensin-1, angiotensin-2 and angiotensin 1-9. Involved in the degradation of atrial natriuretic factor (ANF). Displays UV-

inducible elastase activity toward skin preelastic and elastic fibers.

配列類似性 Belongs to the peptidase M13 family.

細胞内局在 Cell membrane.

画像



All lanes: Anti-CD10 antibody (ab73409) at 1 µg/ml

Lane 1: LNCaP nuclear extract lysate (ab14857)

Lane 2 : Ramos (Human Burkitt's lymphoma cell line) Whole Cell

Lysate

Lane 3: Raji (Human Burkitt's lymphoma cell line) Whole Cell

Lysate

Lane 4: Human kidney tissue lysate - total protein (ab30203)

Lane 5: Human placenta tissue lysate - total protein (ab29745)

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : Goat polyclonal Secondary Antibody to Rabbit IgG - H&L (HRP), pre-adsorbed at 1/5000 dilution

Developed using the ECL technique.

Performed under reducing conditions.


Predicted band size: 85 kDa **Observed band size:** 100 kDa

Additional bands at: 38 kDa. We are unsure as to the identity of

these extra bands.


Exposure time: 2 minutes

CD10 contains a number of potential glycosylation sites (SwissProt) which may explain its migration at a higher molecular weight than predicted.



Immunocytochemistry/ Immunofluorescence - Anti-CD10 antibody (ab73409)

ICC/IF image of ab73409 stained Hek293 cells. The cells were 4% PFA fixed (10 min) and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab73409, 5 μ g/ml) overnight at +4°C. The secondary antibody (green) was **ab96899**, DyLight® 488 goat anti-rabbit lgG (H+L) used at a 1/250 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43 μ M.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-CD10 antibody (ab73409)

IHC image of CD10 staining in Human normal kidney formalin fixed paraffin embedded tissue section, performed on a Leica BondTM system using the standard protocol F. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab73409, 1µg/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors