


# Anti-CD15 antibody [28] ab20137

## 5 References

### 製品の概要

---

製品名	Anti-CD15 antibody [28]
製品の詳細	Mouse monoclonal [28] to CD15
由来種	Mouse
特異性	Specific to CD15. 28 is useful for leukaemia typing and purification of stem cells.
アプリケーション	<b>適用あり:</b> IP, Flow Cyt, IHC-Fr
種交差性	<b>交差が予測される動物種:</b> Human 
免疫原	Tissue, cells or virus corresponding to CD15. Monocytes, separated from other peripheral blood leucocytes (PBL) on fibronectin plates.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

### 製品の特性

---

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	Preservative: 0.02% Sodium azide Constituent: 99.98% PBS
精製度	Protein A/G purified
ポリ/モノ	モノクローナル
クローン名	28
ミエローマ	Sp2/0-Ag14
アイソタイプ	IgM
軽鎖の種類	unknown

## アプリケーション

---

**The Abpromise guarantee**      **Abpromise保証は、次のテスト済みアプリケーションにおけるab20137の使用に適用されず**  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IP		Use at an assay dependent concentration.
Flow Cyt		Use at an assay dependent concentration. <b>ab91545</b> - Mouse monoclonal IgM, is suitable for use as an isotype control with this antibody.
IHC-Fr		Use at an assay dependent concentration. Note: samples should be acetone fixed.

## ターゲット情報

---

<b>関連性</b>	CD15 is a carbohydrate adhesion molecule (and not a protein) that mediates phagocytosis and chemotaxis. Synthesis is directed by FUT4 in lymphoid cells and mature granulocytes, and by FUT9 in promyelocytes and monocytes.
<b>細胞内局在</b>	Golgi Apparatus; Membrane-bound form in trans cisternae of Golgi.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

## Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors