


Anti-CD28 antibody [CD28.2] - Low endotoxin, Azide free ab85986

[11 References](#) [画像数 2](#)

製品の概要

製品名	Anti-CD28 antibody [CD28.2] - Low endotoxin, Azide free
製品の詳細	Mouse monoclonal [CD28.2] to CD28 - Low endotoxin, Azide free
由来種	Mouse
アプリケーション	適用あり: Flow Cyt
種交差性	交差種: Human 交差が予測される動物種: Non human primates 
免疫原	Tissue, cells or virus corresponding to CD28. Human CD28 transfected murine T cell hybridoma.
ポジティブ・コントロール	Flow Cyt: CD28 positive lymphocytes.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term.
バッファー	pH: 7.20 Constituents: 0.14% Monobasic dihydrogen sodium phosphate, 0.87% Sodium chloride
キャリア・フリー	Azide free.
精製度	はい
精製度	Affinity purified
特記事項 (精製)	This purified format is guaranteed to be >90% pure as determined by SDS-PAGE analysis. Endotoxin level is less than or equal to 0.01 EU/ug, as determined by the LaL assay.

ポリ/モノ	モノクローナル
クローン名	CD28.2
アイソタイプ	IgG1

アプリケーション


The Abpromise guarantee **Abpromise保証は、** 次のテスト済みアプリケーションにおけるab85986の使用に適用されます
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
Flow Cyt		Use a concentration of 1 - 12 µg/ml.


ターゲット情報

機能	Involved in T-cell activation, the induction of cell proliferation and cytokine production and promotion of T-cell survival.
組織特異性	Expressed in T-cells and plasma cells, but not in less mature B-cells.
配列類似性	Contains 1 Ig-like V-type (immunoglobulin-like) domain.
細胞内局在	Membrane.

画像



Flow cytometry surface staining pattern of human peripheral blood stained using ab85986 GAM APC.



Separation of human CD28 positive lymphocytes (red-filled) from neutrophil granulocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using ab85986 GAM APC.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors