

Anti-CD44 antibody [EPR1013Y] ab51037

KO 評価済 リコンビナント RabMAb


★★★★☆ 11 Abreviews 153 References 画像数 13

製品の概要

製品名	Anti-CD44 antibody [EPR1013Y]
製品の詳細	Rabbit monoclonal [EPR1013Y] to CD44
由来種	Rabbit
アプリケーション	適用あり: IHC-P, WB 適用なし: Flow Cyt, ICC/IF or IP
種交差性	交差種: Human 非交差種: Mouse, Rat
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	WB: MDA-MB-231, TF-1, HeLa and A549 cell lysates. IHC-P: Pancreatic and cervical cancer, breast and thyroid gland carcinoma, glioma, tonsil and skin tissues.
特記事項	This product is a recombinant monoclonal antibody, which offers several advantages including: <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production For more information see here . Our RabMAb [®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents .

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle.
解離定数 (K _D 値)	K _D = 3.76 x 10 ⁻¹⁰ M



バッファー	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine), 0.5% BSA
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	EPR1013Y
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab51037の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。


アプリケーション	Abreviews	特記事項
IHC-P	★★★★★ (3)	1/50 - 1/100. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
WB	★★★★★ (4)	1/5000. Detects a band of approximately 82 kDa (predicted molecular weight: 82 kDa).

追加情報 Is unsuitable for Flow Cyt, ICC/IF or IP.

ターゲット情報

機能	Receptor for hyaluronic acid (HA). Mediates cell-cell and cell-matrix interactions through its affinity for HA, and possibly also through its affinity for other ligands such as osteopontin, collagens, and matrix metalloproteinases (MMPs). Adhesion with HA plays an important role in cell migration, tumor growth and progression. Also involved in lymphocyte activation, recirculation and homing, and in hematopoiesis. Altered expression or dysfunction causes numerous pathogenic phenotypes. Great protein heterogeneity due to numerous alternative splicing and post-translational modification events.
組織特異性	Isoform 10 (epithelial isoform) is expressed by cells of epithelium and highly expressed by carcinomas. Expression is repressed in neuroblastoma cells.
配列類似性	Contains 1 Link domain.
ドメイン	The lectin-like LINK domain is responsible for hyaluronan binding.
翻訳後修飾	Proteolytically cleaved in the extracellular matrix by specific proteinases (possibly MMPs) in several cell lines and tumors. N-glycosylated. O-glycosylated; contains more-or-less-sulfated chondroitin sulfate glycans, whose number may affect the accessibility of specific proteinases to their cleavage site(s). Phosphorylated; activation of PKC results in the dephosphorylation of Ser-706 (constitutive phosphorylation site), and the phosphorylation of Ser-672.
細胞内局在	Membrane.

画像



All lanes : Anti-CD44 antibody [EPR1013Y] (ab51037) at 1/5000 dilution

Lane 1 : Wild-type HeLa cell lysate

Lane 2 : CD44 knockout HeLa cell lysate

Lane 3 : A549 cell lysate

Lane 4 : LNCaP cell lysate


Lysates/proteins at 20 µg per lane.

Performed under reducing conditions.

Predicted band size: 82 kDa

Observed band size: 75-80 kDa

False colour image of Western blot: Anti-CD44 antibody [EPR1013Y] staining at 1/5000 dilution, shown in green; Mouse anti-Alpha Tubulin [DM1A] ([ab7291](#)) loading control staining at 1/20000 dilution, shown in red. In Western blot, ab51037 was shown to bind specifically to CD44. A band was observed at 75-80 kDa in wild-type HeLa cell lysates with no signal observed at this size in CD44 knockout cell line [ab262515](#) (knockout cell lysate [ab263938](#)). To generate this image, wild-type and CD44 knockout HeLa cell lysates were analysed. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 3 % milk in TBS-0.1 % Tween® 20 (TBS-T) before incubation with primary antibodies overnight at 4 Å°C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed ([ab216773](#)) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed ([ab216776](#)) at 1/20000 dilution.



All lanes : Anti-CD44 antibody [EPR1013Y] (ab51037) at 1/5000 dilution

Lane 1 : Wild-type HeLa (Human epithelial cell line from cervix adenocarcinoma) whole cell lysate

Lane 2 : CD44 knockout HeLa (Human epithelial cell line from cervix adenocarcinoma) whole cell lysate

Lane 3 : A549 (Human lung carcinoma cell line) whole cell lysate

Lane 4 : LNCaP (Human prostate cancer cell line) whole cell lysate

Lysates/proteins at 20 µg per lane.


Performed under reducing conditions.

Predicted band size: 82 kDa


Observed band size: 80 kDa

Lanes 1 -4: Merged signal (red and green). Green - ab51037 observed at 80 kDa. Red - loading control, **ab8245** (Mouse anti-GAPDH antibody [6C5]) observed at 37kDa.

ab51037 was shown to react with CD44 in wild-type HeLa cells in western blot. Loss of signal was observed when CD44 knockout sample was used. Wild-type HeLa and CD44 knockout cell lysates were subjected to SDS-PAGE. Membranes were blocked in 3% milk in TBS-T (0.1% Tween®) before incubation with ab51037 and **ab8245** (Mouse anti-GAPDH antibody [6C5]) overnight at 4°C at a 1 in 5000 dilution and a 1 in 20000 dilution respectively. Blots were incubated with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed (**ab216776**) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD44 antibody [EPR1013Y] (ab51037)



Western blot - Anti-CD44 antibody [EPR1013Y] (ab51037)

Immunohistochemical analysis of paraffin-embedded human pancreatic cancer tissue labeling CD44 with ab51037 at 1/100 dilution followed by goat anti-rabbit IgG H&L (HRP) ([ab97051](#), 1/500). The sample was counter stained with hematoxylin.

Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

All lanes : Anti-CD44 antibody [EPR1013Y] (ab51037) at 1/1000 dilution

Lane 1 : MCF-7 whole cell lysate

Lane 2 : Jurkat whole cell lysate

Lane 3 : MDA-MB-231 whole cell lysate

Lane 4 : HeLa whole cell lysate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution


Predicted band size: 82 kDa

Observed band size: 81 kDa

Exposure time: 3 minutes

Blocking and diluting buffer 5% NFDM/TBST

The expression of CD44 in MCF-7 is low (PMID: 25635866; PMID: 26005723). Jurkat does not express CD44 (PMID: 24127558).



Western blot - Anti-CD44 antibody [EPR1013Y]
(ab51037)


Anti-CD44 antibody [EPR1013Y] (ab51037) at 1/5000 dilution +
TF-1 cell lysate at 10 µg

Secondary

Goat anti-Rabbit HRP labeled. at 1/2000 dilution

Predicted band size: 82 kDa


Observed band size: 82 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-
embedded sections) - Anti-CD44 antibody
[EPR1013Y] (ab51037)

ab51037 (1:100) showing positive staining in human Breast
carcinoma tissue.


Perform heat mediated antigen retrieval with citrate buffer pH 6
before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-
embedded sections) - Anti-CD44 antibody
[EPR1013Y] (ab51037)

Immunohistochemical analysis of paraffin-embedded human tonsil
tissue labeling CD44 with ab51037 at 1/100 dilution followed by
goat anti-rabbit IgG H&L (HRP) ([ab97051](#), 1/500). The sample was
counter stained with hematoxylin.


Perform heat mediated antigen retrieval with citrate buffer pH 6
before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD44 antibody [EPR1013Y] (ab51037)

Immunohistochemical analysis of paraffin-embedded human skin tissue labeling CD44 with ab51037 at 1/100 dilution followed by goat anti-rabbit IgG H&L (HRP) ([ab97051](#), 1/500). The sample was counter stained with hematoxylin.

Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD44 antibody [EPR1013Y] (ab51037)

Immunohistochemical analysis of paraffin-embedded human cervical cancer tissue labeling CD44 with ab51037 at 1/100 dilution followed by goat anti-rabbit IgG H&L (HRP) ([ab97051](#), 1/500). The sample was counter stained with hematoxylin.


Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD44 antibody [EPR1013Y] (ab51037)

ab51037 (1:100) showing positive staining in human Glioma tissue.


Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD44 antibody [EPR1013Y] (ab51037)

ab51037 (1:100) showing positive staining in human Thyroid gland carcinoma tissue.

Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.




OR-D Scanning - Anti-CD44 antibody [EPR1013Y] (ab51037)

Equilibrium disassociation constant (K_D)

Learn more about K_D

[Click here to learn more about \$K_D\$](#)

Why choose a recombinant antibody?



Anti-CD44 antibody [EPR1013Y] (ab51037)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors