abcam

Product datasheet

Anti-Cdc6 antibody [EPR714(2)] ab109315

יולצעבע RabMAb

★★★★★ 2 Abreviews 16 References 画像数7

製品の概要

製品名 Anti-Cdc6 antibody [EPR714(2)]

製品の詳細 Rabbit monoclonal [EPR714(2)] to Cdc6

由来種 Rabbit

アプリケーション 適用あり: ICC/IF, WB, IHC-P

適用なし: Flow Cyt (Intra)

種交差性 交差種: Human

免疫原 Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

ポジティブ・コントロール WB: Hela, 293T, HCT116 treated with hydroxyurea and Raji treated with FBS lysates IHC-P:

Human gastric adenocarcinoma, breast carcinoma, thyroid cancer tissue ICC/IF: HeLa

特記事項 This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility

- Improved sensitivity and specificity

- Long-term security of supply

- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

Mouse: We have preliminary internal testing data to indicate this antibody may not react with this

species. Please contact us for more information.

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long

term. Avoid freeze / thaw cycle.

バッファー pH: 7.20

Preservative: 0.01% Sodium azide

Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine), 0.5% BSA

精製度 Protein A purified

アプリケーション

The Abpromise guarantee <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab109315の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
ICC/IF	*** <u>(2)</u>	1/100. For unpurified use at 1/250 - 1/500
WB		1/1000 - 1/10000. Detects a band of approximately 62 kDa (predicted molecular weight: 63 kDa).
IHC-P		1/90. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol. See IHC antigen retrieval protocols.

追加情報 Is unsuitable for Flow Cyt (Intra).

ターゲット情報


機能 Involved in the initiation of DNA replication. Also participates in checkpoint controls that ensure

DNA replication is completed before mitosis is initiated.

配列類似性 Belongs to the CDC6/cdc18 family.

細胞内局在 Nucleus. Cytoplasm. The protein is nuclear in G1 and cytoplasmic in S-phase cells.

画像



Western blot - Anti-Cdc6 antibody [EPR714(2)] (ab109315)

All lanes : Anti-Cdc6 antibody [EPR714(2)] (ab109315) at 0.5 µg/ml (purified)

Lane 1: 293T (Human embryonic kidney epithelial cell) whole cell

lysates

Lane 2: 293T (Human embryonic kidney epithelial cell) treated with

20mM Hydroxyurea for 24 hours whole cell lysates

Lane 3: HCT116 (Human colorectal carcinoma epithelial cell)

whole cell lysates

Lane 4: HCT116 (Human colorectal carcinoma epithelial cell) treated with 1mM Hydtoxyurea for 16 hours whole cell lysates

Lysates/proteins at 15 µg per lane.


Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) (<u>ab97051</u>) at 1/20000 dilution

Predicted band size: 63 kDa


Blocking and diluting buffer: 5% NFDM/TBST.

Hydroxyurea induces Cdc6 expression (PMID:11346650)




Immunocytochemistry/ Immunofluorescence - Anti-Cdc6 antibody [EPR714(2)] (ab109315)

Immunocytochemistry/ Immunofluorescence analysis of HeLa (Human cervix adenocarcinoma epithelial cell) cells labeling Cdc6 with Purified ab109315 at 1:100 dilution (9.4 μg/ml). Cells were fixed in 4% Paraformaldehyde and permeabilized with 0.1% tritonX-100. Cells were counterstained with Ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor®594) 1:200 (2.5 μg/ml). Goat anti rabbit lgG (Alexa Fluor®488, ab150077) was used as the secondary antibody at 1:1000 (2 μg/ml) dilution. DAPI nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Cdc6 antibody
[EPR714(2)] (ab109315)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human thyroid cancer tissue sections labeling Cdc6 with Purified ab109315 at 1:90 dilution (10.4 µg/ml). Heat mediated antigen retrieval was performed using using ab93684 (Tris/EDTA buffer, pH 9.0)ImmunoHistoProbe one step HRP Polymer (ready to use)was used as the secondary antibody.Negative control:PBS instead of the primary antibody.Hematoxylinwas used as a counterstain




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Cdc6 antibody
[EPR714(2)] (ab109315)

Immunohistochemical analysis of Cdc6 in paraffin embedded Human gastric adenocarcinoma tissue, using unpurified ab109315 at a 1/1000 dilution.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Cdc6 antibody
[EPR714(2)] (ab109315)

Immunohistochemical analysis of Cdc6 in paraffin embedded Human breast carcinoma tissue, using unpurified ab109315 at a 1/1000 dilution.



Immunocytochemistry/ Immunofluorescence - Anti-Cdc6 antibody [EPR714(2)] (ab109315)

This image is courtesy of an abreview by Kirk Mcmanus

Unpurified ab109315 staining Cdc6 in the Hela cell line from Human cervix by ICC/IF

(Immunocytochemistry/immunofluorescence). Cells were fixed with paraformaldehyde, permeabilized with 0.5% Triton X-100. Samples were incubated with primary antibody (1/500 in PBS) for 1 hour at 22°C. Ab150081 (1/200) was used as the secondary antibody.



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors