

Anti-cIAP1 antibody - N-terminal ab196592

1 References [画像数 4](#)

製品の概要

製品名	Anti-cIAP1 antibody - N-terminal
製品の詳細	Rabbit polyclonal to cIAP1 - N-terminal
由来種	Rabbit
アプリケーション	適用あり: IHC-P, WB, ICC/IF
種交差性	交差種: Rat, Human
免疫原	Recombinant fragment corresponding to Human cIAP1 (N terminal). Database link: Q13490
ポジティブ・コントロール	Jurkat, HeLa and HT29 cell extract. Rat kidney tissue.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
バッファー	<p>pH: 7.40</p> <p>Preservative: 0.02% Sodium azide</p> <p>Constituents: 49% PBS, 0.87% Sodium chloride, 50% Glycerol (glycerin, glycerine)</p>
精製度	without Mg ²⁺ and Ca ²⁺
ポリ/モノ	Immunogen affinity purified
アイソタイプ	ポリクローナル
	IgG

アプリケーション


The Abpromise guarantee **Abpromise保証は、** 次のテスト済みアプリケーションにおけるab196592の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-P		1/50 - 1/200.
WB		1/500 - 1/2000. Predicted molecular weight: 70 kDa.
ICC/IF		1/50 - 1/200.

ターゲット情報

機能	Apoptotic suppressor. The BIR motifs region interacts with TNF receptor associated factors 1 and 2 (TRAF1 and TRAF2) to form an heteromeric complex, which is then recruited to the tumor necrosis factor receptor 2 (TNFR2).
組織特異性	Present in many fetal and adult tissues. Mainly expressed in adult skeletal muscle, thymus, testis, ovary, and pancreas, low or absent in brain and peripheral blood leukocytes.
配列類似性	Belongs to the IAP family. Contains 3 BIR repeats. Contains 1 CARD domain. Contains 1 RING-type zinc finger.
細胞内局在	Cytoplasm.

画像




All lanes : Anti-cIAP1 antibody - N-terminal (ab196592) at 1/500 dilution

Lane 1 : Jurkat cell extract

Lane 2 : HeLa cell extract


Lane 3 : HT29 cell extract

Predicted band size: 70 kDa




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-cIAP1 antibody - N-terminal (ab196592)

Immunohistochemical analysis of paraffin-embedded human stomach cancer tissue labeling cIAP1 using ab196592 at a 1/200 dilution.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-cIAP1 antibody - N-terminal (ab196592)

Immunohistochemical analysis of paraffin-embedded rat kidney tissue labeling cIAP1 using ab196592 at a 1/200 dilution.



Immunofluorescence analysis of A549 cells labeling cIAP1 using **ab195692** at a 1/50 dilution.

Immunocytochemistry/ Immunofluorescence - Anti-cIAP1 antibody - N-terminal (ab196592)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors