



Anti-CIKS/ACT1 antibody ab5973

画像数 3

製品の概要

製品名	Anti-CIKS/ACT1 antibody
製品の詳細	Rabbit polyclonal to CIKS/ACT1
由来種	Rabbit
アプリケーション	適用あり: ICC/IF, IHC-P, WB
種交差性	交差種: Human
免疫原	Synthetic peptide corresponding to Human CIKS/ACT1 aa 554-568. Sequence: REEEYVAPPRGPLPT (Peptide available as ab6089 , ab6138) <div>  Run BLAST with  Run BLAST with </div>
ポジティブ・コントロール	WB: Human placental and lung tissue lysates. IHC-P: Humman lung tissue.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle. Stable for 12 months at -20°C.
バッファー	pH: 7.2 Preservative: 0.02% Sodium azide Constituent: PBS
精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション


The Abpromise guarantee Abpromise保証は、 次のテスト済みアプリケーションにおけるab5973の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
ICC/IF		Use a concentration of 20 µg/ml.
IHC-P		Use a concentration of 5 µg/ml.
WB		Use a concentration of 1 µg/ml. Detects a band of approximately 63 kDa (predicted molecular weight: 65 kDa).

ターゲット情報

機能	Could be involved in the activation of both NF-kappa-B via a NF-kappa-B inhibitor kinase (IKK)-dependent mechanism and stress-activated protein kinase (SAPK)/JNK.
組織特異性	Widely expressed.
配列類似性	Contains 1 SEFIR domain.

画像



Western blot analysis of CIKS/ACT1 expression in human placenta tissue lysate with ab5973 at 1 µg /ml. Western blot analysis of CIKS/ACT1 expression in human placenta tissue lysate with ab5973 at 1 µg /ml.



ab5973 at 5µg/ml staining CIKS/ACT1 human lung tissue by IHC.

Immunocytochemistry/ Immunofluorescence - Anti-
CIKS/ACT1 antibody (ab5973)

Immunofluorescence of CIKS/ACT1 in human Lung cells using
ab5973 at 20 µg/ml.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors